

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 31/675, C12N 5/00, 15/00	A1	(11) International Publication Number: WO 94/27612 (43) International Publication Date: 8 December 1994 (08.12.94)
(21) International Application Number: PCT/US94/05548 (22) International Filing Date: 19 May 1994 (19.05.94) (30) Priority Data: 08/063,751 20 May 1993 (20.05.93) US 08/114,222 1 September 1993 (01.09.93) US (71) Applicant: BAYLOR COLLEGE OF MEDICINE [US/US]: One Baylor Plaza, Houston, TX 77030 (US). (72) Inventors: FRENCH, Brent, A.; 5746 Cheena Drive, Houston, TX 77096 (US). RAIZNER, Albert, E.; 11945 N. Durrette, Houston, TX 77024 (US). ROBERTS, Robert; 2198 Troon Road, Houston, TX 77019 (US). (74) Agents: AUERBACH, Jeffrey, I et al.; Howrey & Simon, 1299 Pennsylvania Avenue, NW, Washington, DC 20004 (US).		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: GENETIC THERAPY FOR CARDIOVASCULAR DISEASE (57) Abstract Gene transfer vectors, especially adenoviral vectors, and synthetic vectors capable of emulating specific viral functions, that carry gene sequences capable of ameliorating or preventing the symptoms of cardiovascular disease are disclosed. Such methods can be used to treat restenosis, especially when incident to injury by angioplasty.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

-1-

TITLE OF THE INVENTION:**GENETIC THERAPY FOR CARDIOVASCULAR DISEASE****FIELD OF THE INVENTION:**

5 The invention relates to therapeutic agents for the prevention and treatment of cardiovascular disease. More specifically, the invention relates to modified viruses and/or synthetic complexes capable of emulating specific viral functions. It further relates to the use of such modified viruses in a genetic therapy for
10 cardiovascular disease.

CROSS-REFERENCE TO RELATED APPLICATIONS:

This application is a continuation-in-part of U.S. patent application serial no. 08/063,751, filed on May 20, 1993.

15 **BACKGROUND OF THE INVENTION:**

I. Restenosis and Cardiovascular Disease

Stenosis denotes a narrowing or constriction of a duct or canal. A variety of disease processes, such as atherosclerotic lesions, immunological reactions,
20 congenital deformities, etc., can lead to the stenosis of coronary arteries and thus to myocardial ischemia. Percutaneous transluminal coronary angioplasty (PTCA), the insertion and partial inflation of a balloon catheter into a stenotic vessel to effect its repair,
25 has been extensively used to treat stenosis. In 1990,

-2-

over 300,000 procedures were performed in the United States alone.

The major limitation of PTCA is restenosis (i.e. the re-constriction) of the vascular lesion (Liu, M.W. et al., Circulation 79:1374-1386 (1989), herein incorporated by reference). Restenosis has been found to occur in 30% to 40% of angioplasty patients within 6 months of the procedure (Califf, R.M. et al., J. Amer Col. Cardiol. 17:2B-13B (1991), McBride, W. et al., N. Engl. J. Med. 318:1734-1737 (1988)). Restenosis develops so rapidly that it may be considered a form of accelerated atherosclerosis induced by injury. The significance of the high restenosis rate is compounded by the present inability to predict with a high degree of certainty which patients, vessels, or lesions will undergo restenosis. Indeed, arteries that are widely patent 2 days after PTCA, free of obstructive thrombus, have exhibited restenosis at catheterization 4-6 months later (Liu, M.W. et al., Circulation 79:1374-1386 (1989)).

The mechanism of restenosis is thus not fully understood. The angioplasty procedure causes unavoidable injury to arterial walls. Animal studies of atherosclerosis have suggested that after such balloon injury, denudation of endothelial cells occurs, followed by platelet adhesion and aggregation, and by the release of platelet-derived growth factor (PDGF) as well as other growth factors. The basic biologic factors believed to be involved in the occurrence of restenosis include: the extent of injury, platelets, inflammatory cells, growth factors, cytokines, endothelial cells, smooth muscle cells, and extracellular matrix production.

Both early platelet aggregation and thrombus formation, as well as late myointimal proliferation, are believed to affect the development of the recurrent lesions after PTCA (Liu, M.W. et al., Circulation

-3-

79:1374-1386 (1989)). Although such platelet-thrombus accumulation appears to be a contributing factor in restenosis after balloon dilatation, direct and indirect evidence strongly supports the concept of intimal hyperplasia or proliferation, and migration of vascular smooth muscle cells (VSMC) of medial, or possibly of intimal, origin as the fundamental process of restenosis (see, for review, Liu, M.W. et al., Circulation 79:1374-1386 (1989)). In particular, atherosclerotic coronaries injured by oversized angioplasty balloons, and atherosclerotic coronaries injured by the deployment of intracoronary stents have been found to display vigorous VSMC migration and proliferation (Rodgers, G.P. et al., Circulation 82:560 (1990); Schwartz, R.S. et al., Circulation 82:2190 (1990); Karas, S.P. et al., J. Amer. Col. Cardiol. 20:467 (1992); Johnson, D.E. et al., Circulation 78(suppl. II):II-82 (1988), all herein incorporated by reference). There is also a large body of indirect and experimental data. Intimal thickening induced by balloon injury in normal rat arteries reaches its maximum at 2 months, a time course of the intimal growth that approximates that observed in humans after PTCA (Serruys, P.W. et al., Circulation 77:361-371 (1988)). Intimal hyperplasia is also known to be an important component of restenosis in the atherosclerotic rabbit artery. Medial smooth muscle cells are the major cellular component of the arterial wall, and the intimal proliferation of these cells is the only major reparative or reactive response of the arterial wall to mechanical or inflammatory injury.

In their normal quiescent state, smooth muscle cells proliferate at a very low rate (probably less than 0.1% per day), in response to the presence of growth inhibitory factors, such as heparin. In contrast, endothelial cells, macrophages, and platelets are thought to provide the mitogenic stimulus necessary to the growth of medial smooth muscle cells in normal

vessels (Liu, M.W. *et al.*, Circulation 79:1374-1386 (1989)). It has been proposed that situations such as vessel damage cause platelets to aggregate on the angioplasty-induced wound surface. These activated platelets release substances that promote local vasoconstriction and thrombus formation and growth factors that activate mesenchymal cells in the vicinity of injured tissue. Within a few hours monocytes also appear, increasing in number over the first few days. Like platelets, monocytes secrete growth factors capable of initiating and promoting local tissue mesenchymal cell migration. Such activated platelets and cells produce and release a number of growth stimulating factors (such as platelet derived growth factor (PDGF), epidermal growth factors, insulin-like growth factors, transforming growth factors (TGF- α and TGF- β), PDGF-like molecules, fibroblast growth factors (acidic and basic), interleukin-1 and somatomedin-C, etc.) that are capable of inducing muscle cell proliferation (Liu, M.W. *et al.*, Circulation 79:1374-1386 (1989); Forrester, J.S. *et al.*, J. Amer. Col. Cardiol. 17:758-769 (1991); Stiles, C.D. *et al.*, J. Supramolec. Struct. 13:489-499 (1980)). In this manner, PDGF and other growth factors are released from platelets and other cells to assist in the repair process.

Studies examining the proliferative activity of smooth muscle cells have found that most enter the growth cycle between 2 and 3 days after balloon injury and the vast majority of proliferation is completed within 7 days Clowes, A.W. *et al.*, Circ. Res. 56:139-145 (1985)). In addition, the population of nondividing smooth muscle cells remains relatively constant between 7 and 14 days, suggesting that if smooth muscle cells proliferate, they do so soon after injury. Smooth muscle cell proliferation thus appears to be an acute response related to the initial injury.

-5-

II. Treatment of Restenosis

Mechanical approaches aimed at minimizing or preventing restenosis with atherectomy devices, stents, or specialized balloon catheters have not, to date, significantly reduced the restenosis rate (Waller, B.F., J. Amer Col. Cardiol. 13:969-987 (1989)). Systemic drug therapy has yet to be effective (Chesebro, J.H. et al., Circulation 80:II-64 (1989); Pepine, C.J. et al., Circulation 81:1753-1761 (1990); Liu, M.W. et al., Circulation 79:1374-1386 (1989), all herein incorporated by reference)).

Intracoronary site specific gene transfer has been proposed as a means for allowing the production of therapeutic proteins in concentrations sufficient to combat restenosis (Swain, J.L., Circulation 80:1495-1496 (1989); O'Brien, T.X. et al., Circulation 83:2133-2136 (1991)). The general principles of gene therapy have been discussed by Oldham, R.K. (In: Principles of Biotherapy, Raven Press, NY, 1987); Boggs, S.S. (Int. J. Cell Clon. 8:80-96 (1990)); Karson, E.M. (Biol. Reprod. 42:39-49 (1990)); Ledley, F.D., In: Biotechnology, A Comprehensive Treatise, volume 7B, Gene Technology, VCH Publishers, Inc. NY, pp 399-458 (1989)); all of which references are incorporated herein by reference.

Despite the potential for intracoronary site specific gene transfer to allow production of therapeutic proteins in concentrations sufficient to combat restenosis, methods involving retroviruses and other transfection approaches have proven to be unsuccessful. Wilson, J.M. et al. (Science 244:1344-1346 (1989)) and Nabel, E.G. et al. (Science 244:1342-1343 (1989)) have discussed the use of retroviruses to introduce reporter genes into endothelial cells in vitro. The genetically altered cells were reintroduced into vascular segments of intact animals and found to be able to express the marker protein (β -galactosidase) for

-6-

at least 5 weeks. In an in vitro study, Dichek, D.A. et al. (Circulation 80:1347-1353 (1989)) discussed the ability of cDNA-transfected sheep endothelial cells that had been seeded onto metallic stents to produce

5 physiologically active tissue plasminogen activator.

Nabel, E.G. et al. (Science 249:1285-1288 (1990)) discuss the use of a dual-balloon catheter system to directly transfect pig iliofemoral arteries and bypass the steps of in vitro transfection and reimplantation of

10 endothelial cells. Both retroviruses and cationic liposomes were used as vectors. In order to obtain transfection, these investigators incubated femoral artery segments for 30 minutes with DNA-Lipofectin complexes delivered by means of a dual-balloon catheter

15 that did not provide for perfusion to the distal vasculature. This method proved effective in peripheral vessels, but perfusion of blood to the distal artery would be necessary during coronary transfection in order to avoid cardiac ischemia, arrhythmia or infarction

20 (Chapman, G.D. et al., Circ. Res. 71:27-33 (1992)).

As indicated, attempts have been made to use cationic liposomes as the transfection vehicle (Nabel, E.G. et al., Science 249:1285-1288 (1990); Lim, C.S. et al., Circulation 83:2007-2011 (1991)). Lim, C.S. et al.

25 (Circulation 83:2007-2011 (1991)) discuss the direct transfection in vivo of the coronary arteries of an intact canine by a surgical method using cationic liposomes. A recent report by Chapman, G.D. et al. (Circ. Res. 71:27-33 (1992)), discusses the possibility

30 of using Lipofectin™-mediated transfection of coronary arteries. As indicated in that report, transfection was extremely inefficient.

Although various researchers have reported in vitro methods for gene transfer, extrapolating from such in

35 vitro work to in vivo intracoronary gene transfer with a catheter technique poses special problems. A prolonged incubation of the gene of interest in the coronary

-7-

artery is desirable, since in vitro expression of protein is directly correlated with the duration of exposure of cells to DNA (Chapman, G.D. et al., Circ. Res. 71:27-33 (1992); Lim, C.S. et al., Circulation 83:2007-2011 (1991)).

Thus, whereas direct in vivo gene transfer into peripheral arteries has been demonstrated using Lipofectin™ or retroviral vectors (Nabel, E.G. et al., Science 249:1285-1288 (1990)), and canine coronary arteries have been targeted with Lipofectin™ (Lim, C.S. et al., Circulation 83:2007-2011 (1991); Chapman, G.D. et al., Circ. Res. 71:27-33 (1992)), the low efficiencies inherent in these gene transfer methods have precluded clinical application.

Thus, the application of gene therapy has been limited by the low efficiencies of gene transfer afforded by conventional and recent methods. In particular, a more efficient technique of gene transfer that would be applicable not only to normal coronary arteries, but also to atherosclerotic vessels, and particularly those displaying the histopathology of restenosis would be highly desirable. The present invention provides such a therapy.

BRIEF DESCRIPTION OF THE FIGURES:

Figure 1 shows a representation of the construction of an adenoviral vector carrying the luciferase gene. The small plasmid at right carries the luciferase reporter gene embedded in sequences from the left end of the adenovirus type 5 genome, while the larger plasmid at left carries the complete dl309 adenovirus type 5 genome. The "x" between the two plasmids represents the homologous crossover event which generates the desired recombinant virus following the cotransfection of permissive 293 host cells.

-8-

Figure 2 shows a comparison of Lipofectin™- and adenovirus-mediated gene transfer into normal, atherosclerotic ("Athero") and restenotic porcine coronary arteries ("Ballooned" and "Stented"). The mean picograms (pg) of active luciferase recovered per coronary artery undergoing each of the eight regimens are indicated on the y-axis. Error bars indicate standard deviation of the mean.

Figure 3 shows three graphs demonstrating the levels of gene expression resulting in seven neonatal rat tissues following the direct injection of: i) Ad5/RSV/GL2 into the carotid artery, ii) Ad5/sk394/Luc into the carotid artery, and iii) Ad5/RSV/GL2 into the peritoneal cavity.

SUMMARY OF THE INVENTION:

The invention concerns therapeutic agents for the prevention and treatment of cardiovascular disease. More specifically, the invention relates to modified viruses, especially adenoviruses, that carry gene sequences capable of ameliorating or preventing the symptoms of cardiovascular disease. It further relates to the use of such modified viruses in a genetic therapy for cardiovascular disease.

In detail, the invention provides a method of genetic therapy for the treatment of cardiovascular disease, wherein the method comprises providing an effective amount of a gene transfer vector to a somatic cell of an animal, the vector comprising a therapeutic gene sequence and being sufficient to mediate efficient gene transfer.

The invention further concerns the embodiment wherein the gene transfer vector is derived from adenovirus Subgenus C.

The invention further concerns the embodiment wherein the gene transfer vector is a complex derived

-9-

from two or more of the following components or their derivatives: DNA, protein, cationic polymer, cationic lipid, and/or synthetic molecules, and in particular, the sub-embodiment wherein the protein-based and/or synthetic molecules of the gene transfer vector provide at least one of the following functions: i) plasma membrane penetration (by membrane fusion or endocytosis), ii) escape from lysosomal degradation (preferably by endosomal avoidance or disruption), iii) nuclear transport (preferably by virtue of a nuclear localization sequence), and iv) persistent nuclear retention (preferably by resistance to nuclease activity or by providing for replication).

The invention further concerns the above embodiments wherein the structure of the complex is maintained by ionic interactions between the anionic DNA-based component and (cationic polymers and/or cationic lipids) or wherein the proteins and/or synthetic molecules are chemically conjugated with a cationic polymer to maintain interaction with the DNA-based component or wherein the proteins and/or synthetic molecules are incorporated into liposomes (or liposomal complexes) of cationic lipids in order to maintain association with the DNA-based component.

The invention particularly concerns the embodiments, wherein the vascular cell is a smooth muscle cell, an endothelial cell, an immune cell, a cardiomyocyte or a cardiac fibroblast or wherein the cardiovascular disease is restenosis, atherosclerosis or dyslipidemia or involves inflammation, thrombosis, ischemia, myocardial infarction, reperfusion injury, or the immunological rejection of transplanted organs or tissues.

The invention also concerns the embodiment wherein the therapeutic gene sequence is expressed by the vascular cell, or secreted by the vascular cell, or is arrayed as a cell surface receptor by the vascular cell,

-10-

and particularly wherein the therapeutic gene sequence encodes a receptor, receptor antagonist, a soluble receptor, or a receptor protein, containing a dominant-negative mutation.

5 The invention also concerns the embodiment wherein the therapeutic gene sequence is selected from the group consisting of Thrombin inhibitors (factor Xa antagonists, hirudin, thrombomodulin, antithrombin III, protein C, protein S, lipoprotein-associated coagulation
10 inhibitor), Platelet inhibitors (RGD peptides, GP IIb/IIIa receptor antagonists), GF inhibitors (calcitonin gene-related peptide, somatostatin, soluble growth factor receptors, dominant-negative receptors, receptor antagonists), Antioxidants (superoxide
15 dismutase, catalase, glutathione peroxidase), Antiproliferative proteins (tyrosine phosphatase, γ interferon, p53, eIF-2 kinase), Chimeric toxins (FGF-saporin, EGF-diphtheria toxin, TGF α -pseudomonas exotoxin A), Anti-inflammatory agents, TGF β , lipocortin,
20 vasocortin, IL-10), Anti-lipemic agents (LDL receptor, ApoA1, phospholipases) and Antisense or Ribozyme RNAs targeted against: c-myc, c-myb, c-fos, c-jun, PCNA, tyrosine kinase receptors, nucleolar p120 protein, slow calcium channels, HMG-CoA reductase, cholesterol
25 acyltransferase, cholesterol ester transfer proteins, thromboxane synthase, thromboxane receptors, etc.

 The invention also contemplates the embodiments wherein the effective amount of the vector is provided in vivo by a percutaneous, catheter-based delivery, or
30 by direct intra-arterial injection or infusion, with or without surgical isolation of the vessel, or is provided ex vivo to a graft, and wherein the graft is then introduced into the animal.

 The method can be performed prior to,
35 simultaneously with or subsequent to angioplasty, or surgical procedure, and may comprise the administration of more than one therapeutic gene sequences to the

-11-

animal. The method may additionally comprise administering to the animal an additional pharmacological agent(s) in conjunction with gene therapy.

5 **DESCRIPTION OF THE PREFERRED EMBODIMENTS:**

I. Genetic Therapy for Cardiovascular Disease

 The present invention concerns the use of viral
vectors to provide a gene therapy for cardiovascular
disease. In particular, it provides vectors and methods
10 for transferring genetic material into the somatic cells
of animals, including humans in order to effect such
genetic therapy. The somatic cells that are the
recipients of the gene therapy include smooth muscle
cells, cardiomyocytes, cardiac fibroblasts, endothelial
15 cells (such as the cells that line the vasculature),
hepatocytes, neuronal cells, and cells of the immune
system (such as stem cells, leukocytes, etc.).

 The present invention derives in part from the
recognition that despite the fact that clinically useful
20 gene therapy protocols for treating cardiovascular
disease have not heretofore been developed, such
diseases are particularly amenable to gene therapy
since: (i) genes can be transferred directly into
coronary arteries, (ii) a number of proteins have been
25 shown to inhibit VSMC proliferation, and (iii) even a
temporary inhibition of VSMC proliferation might be
sufficient to limit the formation of restenotic lesions
(Forrester, J.S. et al., J. Amer. Col. Cardiol. 17:758-
769 (1991)). In particular, it is desirable to be able
30 to deliver genes directly into any of four types of
coronary vessels: (i) normal coronaries, (ii) athero-
sclerotic coronaries, (iii) atherosclerotic coronaries
injured by angioplasty balloons, and (iv) athero-

-12-

sclerotic coronaries in which intra-coronary stents have been deployed.

The "treatment" provided by the molecules and methods of the present invention may be either

5 "prophylactic" or "therapeutic." A prophylactic treatment is one that is provided in advance of any symptom of cardiovascular disease in order to prevent or attenuate any subsequent onset of a symptom of such disease. A therapeutic treatment is one that is

10 provided in response to the onset of a symptom of cardiovascular disease, and serves to attenuate an actual symptom of the disease.

As used herein, the term "disease" denotes genetic or acquired disease, as well as conditions or disorders

15 that may or may not have a genetic component (such as, for example, myocardial infarction, aneurysm, atherosclerosis, etc.). In particular, the term "disease" includes trauma or injury that is incidental to surgery, or other treatment, such as incidental to

20 angioplasty, or to the deployment of intracoronary stents, etc.

Any of an array of cardiovascular diseases are amenable to the gene therapy protocols of the present invention. One particularly preferred class of

25 cardiovascular diseases and conditions that may be treated by the methods of the present invention is associated with a process of cellular migration. Such migration may, for example, comprise an inflammatory response, such as the migration of neutrophils,

30 lymphocytes and other leukocytes to sites of inflammation. Alternatively, such migration may involve the movement of muscle cells to sites of injury as is observed in atherosclerosis and restenosis (either spontaneous, incident to myocardial infarction, or

35 incident to surgical intervention, such as angioplasty). A second preferred class of diseases and conditions that can be treated by the methods of the present invention

-13-

involve those that are associated with the endogenous expression of hormonal and other factors, such as growth hormone, growth factors, etc.

5 The present invention can therefore be employed in the treatment of hyperlipidemia (such as high cholesterol levels, etc.), hereditary myopathies, thrombosis, restenosis, ischemic heart disease, atherosclerosis, asthma; adult respiratory distress syndrome (ARDS) or multiple organ injury syndromes
10 secondary to septicemia or trauma; reperfusion injury of myocardial or other tissues; acute glomerulonephritis; reactive arthritis; dermatoses with acute inflammatory components; acute purulent meningitis or other central nervous system inflammatory disorders; thermal injury;
15 hemodialysis; leukapheresis; ulcerative colitis; Crohn's disease; necrotizing enterocolitis; granulocyte transfusion associated syndromes; and cytokine-induced toxicity.

II. The Molecules of the Present Invention

20 A. The Therapeutic Gene Sequences of the Present Invention

The invention accomplishes the treatment of such diseases by providing vectors and methods suitable for delivering "therapeutic gene sequences" to the somatic
25 cells of patients. The particular therapeutic gene sequence used will vary depending on the condition that is to be treated, and the targeted somatic cell type. Thus, for example, muscle cells can be targeted with gene sequences that effect their ability to support the
30 migration of smooth muscle cells in order to treat potential restenosis, etc., or leukocytes in order to treat inflammatory responses, etc. Neuronal cells can be targeted to receive sequences that would relieve cardiac conduction abnormalities.

-14-

The therapeutic gene sequences of the present invention can, and preferably will, encode therapeutic proteins and antisense RNAs and/or ribozymes. In an especially preferred sub-embodiment, the therapeutic gene sequences will direct the secretion of the therapeutic protein. In an alternative embodiment, the therapeutic gene sequences of the present invention may provide antisense RNA (European Patent Appln. Publ. Nos. 263,740; 335,451; and 329,882, PCT Appln. Publ. No. WO90/00624) or ribozyme to the target cell in order to repress or prevent the expression of an endogenous gene sequence.

The transcription of the therapeutic gene sequences of the invention can be mediated from any suitable eucaryotic promoter. Examples of such suitable promoters include the RSV LTR, the SV40 early promoter, the cytomegalovirus (CMV) IE promoter, and the MMTV promoter. The SV40 promoter appears to be less active than the RSV LTR following transfection of cultured coronary VSMC. The CMV promoter appears to be stronger than the RSV LTR. Analysis of a luciferase vector driven by the smooth muscle α -actin promoter has suggested that it is nearly inactive in cultured coronary VSMC.

The nature of the therapeutic gene sequence employed, and whether that sequence is to be expressed and/or secreted will depend upon the particular disease or condition that is to be treated. As indicated, the expression of these molecules can be depressed by antisense expression, however, it is more preferable to employ therapeutic gene sequences that direct the secretion of proteins that are antagonists of inflammatory mediators or proliferation or migration processes. Examples of such therapeutic gene sequences are those that encode Thrombin inhibitors (factor Xa antagonists, hirudin, thrombomodulin, antithrombin III, protein C, protein S, lipoprotein-associated coagulation

-15-

inhibitor), Platelet inhibitors (RGD peptides, GP IIb/IIIa receptor antagonists), GF inhibitors (calcitonin gene-related peptide, somatostatin, soluble growth factor receptors, dominant-negative receptors, receptor antagonists), Antioxidants (superoxide dismutase, catalase, glutathione peroxidase), Antiproliferative proteins (tyrosine phosphatase, γ -interferon, p53, elF-2 kinase), Chimeric toxins (FGF-saporin, EGF-diphtheria toxin, TGF α -pseudomonas exotoxin A), Anti-inflammatory agents, TGF β , lipocortin, vasocortin, IL-10), Anti-lipemic agents (LDL receptor, ApoA1, phospholipases) and Antisense or Ribozyme RNAs targeted against: c-myc, c-myb, c-fos, c-jun, PCNA, tyrosine kinase receptors, nucleolar p120 protein, slow calcium channels, HMG-CoA reductase, cholesterol acyltransferase, cholesterol ester transfer proteins, thromboxane synthase, thromboxane receptors, etc. The strategy behind this approach derives from the fact that only a subset of the vascular cells will express therapeutic genes after gene transfer. In order to facilitate the proper processing of the natural propeptide sequences by the vascular smooth muscle cells, it may be desirable to link the therapeutic gene sequences to gene sequences encoding signal peptides known to be efficiently processed by vascular cells (such as those from collagen or lysyl oxidase).

The secretion of therapeutic gene products from a modest population of infected vascular cells will create a microenvironment around the lesion which contains a high concentration of antiproliferative proteins. For example, locally-produced, persistent high concentrations of γ -interferon, somatostatin, etc., will inhibit VSMC proliferation and thus prevent or lessen restenosis.

In a second sub-embodiment, the therapeutic gene sequences will direct the secretion of the therapeutic protein that is a "soluble receptor" (i.e. a fragment of

-16-

a receptor molecule that retains its ligand binding capacity, but is secreted from the cell into the circulation, or intercellular fluid). This approach, referred to as a "soluble receptor strategy," will enable a relatively small number of genetically reprogrammed cells to protect their neighbors from proliferative stimuli by secreting soluble receptors which will bind and thus inactivate specific growth factors. Examples of suitable receptors that may be solubilized include the PDGF α - and β -receptors, the TGF α - and β -receptors, receptors for the acidic and basic forms of FGF, EGF receptors, thrombin receptors, interleukin receptors, macrophage colony stimulating factor receptors, and TNF- α receptors, as well as soluble ICAMS, VCAMS, soluble selectins, etc.

In a third sub-embodiment, the therapeutic gene sequences will direct the expression of the therapeutic protein and its insertion into the target cell membrane, so as to create a receptor molecule on the target cell that would be capable of contacting extracellular molecules or ligands. These recombinant receptors may be used to increase cellular responsiveness to endogenous growth inhibitors. Alternatively, these recombinant receptors may be designed so as to transmit a growth regulatory signal in response to the administration of an exogenous substance. In this manner, the neointimal growth rate can be precisely controlled by systemic administration of the exogenous substance.

In a fourth sub-embodiment, the therapeutic gene sequences will direct the expression of growth factor receptors which have been mutated or truncated so that they are no longer capable of signal transduction. For example, the deletion or mutation of the tyrosine kinase domain in a growth factor receptor cDNA will result in a protein which is capable of dimerization (or multimerization) and ligand binding; however, ligand

-17-

binding will not lead to signal transduction due to the lack of a functional kinase domain. Defective receptors that have such characteristics are referred to as dominant-negative receptors, because the incorporation of one mutant receptor can inactivate a larger multimeric complex, so that the negative effect of the mutant receptor can dominate over the wild-type binding partner(s).

1. Adenovirus-mediated Gene Therapy for Atherosclerosis

The preferred gene therapy for atherosclerosis derives in part from an appreciation of the molecular machinery for lipid metabolism and the existence of single-gene disorders such as LDL receptor deficiency that are especially propitious candidates for gene therapy in somatic cells. Factors which favor LDL receptor deficiency as a priority for gene therapy research include extraordinarily efficient uptake of adenovirus by liver even with systemic injection, the opportunity for portal vein injection of virus or engineered cells, the likelihood that relatively broad ranges in the levels of gene expression will be tolerated, the early lethality resulting from LDL receptor mutations, and the potential implications for augmenting LDL receptor abundance in more common forms of hyperlipidemia.

A reciprocal strategy to protect against or reverse atherosclerosis is based on augmentation of circulating HDL cholesterol: in mice, intravenous injection of recombinant adenovirus encoding human apolipoprotein A1 results in overexpression of ApoA1 and, consequently, increasing HDL cholesterol to levels that are known to be protective in humans. ApoA1, like other secreted, circulating proteins, is particularly amenable to a gene transfer strategy, since virtual any somatic cell

-18-

(particularly hepatic cells) could be targeted to serve as a source of the gene product.

2. Adenovirus-mediated Gene Therapy for Hereditary Myopathies

5 Gene delivery to muscle itself will be required for
Duchenne (DMD) and Becker muscular dystrophies: X-linked
disorders caused by absence or abnormalities,
respectively, of the sarcolemmal protein, dystrophin.
Myoblast transfer and direct injection of plasmids
10 encoding dystrophin have been employed in efforts to
restore dystrophin expression in DMD patients or
dystrophin-deficient mdx mice, yet daunting limitations
are inherent to both techniques that rely on local
injection, i.e. the total mass of skeletal muscle
15 needing correction, the distribution and inaccessibility
of respiratory muscles on which the clinical outcome
largely depends, the concurrent need to restore
dystrophin expression in myocardium, and the meager
production of dystrophin-positive muscle fibers.

20 In contrast, mouse skeletal muscle that had been
injected with adenovirus encoding the E. coli β -
galactosidase marker gene (lacZ) labeled more extensive
muscle segments than did plasmid injection. Indeed,
adenovirus was at least 10-fold more efficient than
25 naked DNA for conferring expression of dystrophin to mdx
mice. Of note, however, some therapeutic gene
sequences, such as dystrophin are quite large (14,000
bp) and substantially exceed the present (7,500 bp)
capacity of adenovirus vectors. Thus, for such
30 therapies, truncated molecules (such as the 6,300 bp
"minidystrophin" cDNA, derived from a Becker patient
with phenotypically mild deletion are preferably
employed. In this circumstance, even the truncated
protein was sufficient for appropriate in vivo
35 localization. It is possible that patients suffering

-19-

from more severe deletions may not fully respond to adenoviral delivery of minidystrophin.

As demonstrated in Example 8 below, cardiac muscle can be targeted at high efficiency and specificity by simple intra-arterial injection. Skeletal muscle can also be targeted at lower efficiency and specificity by simple intraperitoneal injection. Nonetheless, muscle-specific regulatory sequences and even elements that discriminate between cardiac and skeletal muscle may be used, to direct tissue-specific expression of recombinant adenoviral genes in clinical settings where local delivery would be problematic.

3. Adenovirus-mediated Gene Therapy for Coronary Artery Restenosis and Thrombosis

As indicated above, adenovirus-mediated gene therapy is particularly well-suited as an adjunct to coronary angioplasty, since even temporary inhibition of smooth muscle cell proliferation might suffice to limit formation of restenotic lesions that confound the clinical outcome of this procedure. Furthermore, although numerous pharmacological agents inhibit restenosis in rodent models, these approaches have largely failed when attempted in large mammals or in clinical trials. As this failure could be due in part to the inability of large mammals (including humans) to tolerate the high systemic doses used in rodent models, local delivery catheters might be employed to deliver high-dose therapy locally to the susceptible coronary artery in conjunction with cardiac catheterization or angioplasty. However, many recombinant pharmaceuticals are prohibitively expensive for chronic treatment, and there are formidable technical challenges to maintaining high local drug doses in coronary vessels for days to weeks after a single transcatheter infusion. Gene therapy, by contrast, is particularly well suited to

these challenges, since this approach would generally reprogram the coronary vessel, to produce the necessary recombinant proteins for the requisite period of time. As indicated below, such gene therapy can be employed in concert with the above-discussed pharmacological treatments.

The concept of direct gene transfer to inhibit restenosis was first articulated in the context of retrovirus- and Lipofectin™-mediated gene transfer (Nabel, E.G. *et al.*, Science 249:1285 (1990)). However, the absolute level of recombinant protein produced *in vivo* by these techniques was far below that which would be considered therapeutically significant.

Apart from the need for more efficient gene transfer, methods were required that would be applicable to atherosclerotic vessels as well. The potential efficiency for adenovirus-mediated gene transfer to the vasculature *in vivo* is supported strongly by *in vitro* cell culture studies: coronary smooth muscle cells of canine, porcine, and human origin have been infected with efficiencies approaching 100%. Nanogram amounts of recombinant gene product can be recovered from a single 35 mm cell culture plate as early as 24 hours following infection. Successful gene transfer into virtually each cell on a culture plate represents a remarkable improvement over conventional transfection techniques (calcium-phosphate precipitation, retroviral transduction, Lipofectin™, DEAE-dextran, electroporation), which typically modify fewer than 10% of treated cells.

Recombinant adenovirus is an efficient vector for *in vivo* gene transfer and can be preferentially directed at vascular endothelium or smooth muscle cells. Recombinant adenoviral vectors have also been applied to the peripheral vessels of sheep (Lemarchand, P. *et al.*, Circ. Res. 72:1132-1138 (1993)) and rabbits (Willard, J.E., *et al.*, Circulation 86:I-473 (1992)), targeting

-21-

the endothelium and intimal smooth muscle, respectively, in intact or denuded vessels. Trans-catheter delivery has been shown to be feasible for targeting adenoviral vectors to rabbit coronary arteries and even myocardium (Barr, E. et al., J. Cell. Biochem. 17D:195 (1993), herein incorporated by reference).

Suitable sources of therapeutic cDNA or gene sequences are abundant in the literature. Thrombin inhibitors (factor Xa antagonists (Dunwiddie, C.T. et al., Biochem. 31:12126-12131 (1992)), hirudin (Degryse, E. et al., Prot. Engineer. 2:459-465 (1989)), thrombomodulin (Wen, D.Z. et al., Biochem. 26:4350-4357 (1987)), antithrombin III (Gillespie, L.S. et al., J. Biol. Chem. 266:3995-4001 (1991)), protein C (Foster, D.C. et al., Biochem. 26:7003-7011 (1987)), protein S (Chang, G.T. et al., Thromb. Haemost. 67:526-532 (1992)), lipoprotein-associated coagulation inhibitor (Girard, T.J. et al., Thromb. Res. 55:37-50 (1989)), Platelet inhibitors (RGD peptides (Dennis, M.S. et al., Proteins 15:312-321 (1993)), GP IIb/IIIa receptor antagonists (Chang, H.H. et al., Biochem. Biophys. Res. Commun. 190:242-249 (1993)), GF inhibitors (calcitonin gene-related peptide (Noel, M. et al., J. Endocrinol. Invest. 13:567-573 (1990)), somatostatin (Shen, L.P. et al., Proc. Natl. Acad. Sci. (U.S.A.) 79:4575-4579 (1982)), soluble growth factor receptors (Duan, D.S. et al., J. Biol. Chem. 266:413-418 (1991)), dominant-negative receptors (Amaya, E. et al., Cell 66:257-270 (1991)), receptor antagonists (Baird, A. et al., Proc. Natl. Acad. Sci. (U.S.A.) 85:2324-2328 (1988)), Antioxidants (superoxide dismutase (Ho, Y.-S. et al., FEBS Lett. 229:256-260 (1988)), catalase (Furuta, S. et al., J. Biochem (Tok) 107:708-713 (1990)), glutathione peroxidase (Ho, Y.-S. et al., FEBS Lett. 301:5-9 (1992)), Antiproliferative proteins (tyrosine phosphatase (Gu, M. et al., Proc. Natl. Acad. Sci. (U.S.A.) 89:2980-2984 (1992)), γ -interferon (Devos, R.

- et al., Nucl. Acid. Res. 10:2487-2501 (1982)), p53
 (Zakut-Houri, R. et al., EMBO J. 4:1251-1255 (1985)),
 eIF-2 kinase (Meurs, E. et al., Cell 62:379-390 (1990)),
 Chimeric toxins (FGF-saporin (Lappi, D.A. et al.,
 5 Biochem. Biophys. Res. Commun. 160:917-923 (1989)), EGF-
 diphtheria toxin (Pickering, J.G. et al., J.
Clin. Invest. 91:724-729 (1993)), TGF α -pseudomonas
 exotoxin A (Mesri, E.A. et al., J. Biol. Chem. 268:4853-
 4862 (1993)), Anti-inflammatory agents (TGF β (Bourdrel,
 10 L. et al., Prot. Express. Purif. 4:130-140 (1993)),
 lipocortin (Frey, B.M. et al., Biochem J. 275:219-225
 (1991)), vasocortin (Sautebin, L. et al., Pharmacol. Res.
25:1-12 (1992)), IL-10 (Goodman, R.E. et al., Biochem.
Biophys. Res. Commun. 189:1-7 (1992)), Anti-lipidemic
 15 agents (LDL receptor (Herz, J. et al., Proc. Natl. Acad.
Sci. (U.S.A.) 90:2812-2816 (1993)), ApoA1 (Kopfler, W.P.
 et al., Clin. Res. 41:P211A (1993)), phospholipases
 (Kanda, A. et al., Biochim. Biophys. Acta. 1171:1-10
 (1992)) and antisense or ribozyme RNAs (targeted
 20 against: c-myc, c-myb (Sala, A. et al., Proc. Natl.
Acad. Sci. (U.S.A.) 89:10415-10419 (1992)), c-fos or c-
 jun, PCNA (Smith, M.J. et al., Blood 79:2107-2115
 (1992)), tyrosine kinase receptors, nucleolar p120
 protein (Saijo, Y. et al., Canc. Lett. 68:95-104
 25 (1993)), slow calcium channels, thrombospondin receptors
 (Silverstein, R.L. et al., J. Biol. Chem. 267:16607-
 16612 (1992)), HMG-CoA reductase, cholesterol
 acyltransferase, cholesterol ester transfer proteins,
 thromboxane synthase, thromboxane receptors, etc.).
 30 These therapeutic cDNAs can be incorporated into
 replication-deficient adenoviral vectors by the methods
 outlined in Section IIIB, Example 2 and diagrammed in
 Figure 1. In one embodiment, molecular cloning methods
 are used to replace the luciferase cDNA in
 35 pXCJL.1/RSV/GL2 with the therapeutic cDNA to create an
 adenoviral shuttle vector carrying the therapeutic cDNA
 between the RSV promoter and the SV40 polyadenylation

-23-

5 signal. Cotransfection of the 293 host cell line with this construction and the balance of the adenoviral genome (provided by pJM17) will produce the desired therapeutic virus by homologous recombination. Standard protocols found in published procedures can be used to isolate, plaque purify, and amplify the recombinant virus.

10 The procedure described above is intended to illustrate, but not to limit the present invention. For example, a proven expression cassette consisting of a minimum of a promoter, a therapeutic cDNA, and a polyadenylation signal can substitute for the luciferase expression cassette in pXCJL1/RSV/GL2, as any combination of 5'-end, therapeutic cDNA (or gene), and 15 3'-end are possible. Furthermore, alternative adenoviral cloning systems can be employed in which the therapeutic expression cassette is inserted elsewhere in the adenoviral genome (such as the E3 region). The replication-deficiency of such a virus would be 20 maintained by the E1 deletion. Although the example detailed above describes expression cassettes which are less than about 5.5 kilobases long, longer inserts can be accommodated by deleting additional regions in the adenoviral genome. For example, deletion of both the E1 25 and E3 regions will accommodate inserts of up to about 7.5 kilobases in length. Insertions of up to 35 kilobases can be accommodated by adding the adenoviral inverted terminal repeats (ITRs) to the desired DNA fragment, then transfecting a host cell line which 30 provides additional viral functions in trans, or transfecting host cells in the presence of a helper virus.

35 An additional gene therapy can be applied following difficult angioplasty cases (involving coronary dissection or thrombosis-prone arterial segments), cases of emergency (rescue) angioplasty, and other clinical scenarios where coronary thrombosis might pose a chronic

threat. The threat of thrombosis could be substantially reduced for a period of weeks by including a procedure whereby an anti-thrombotic gene therapy (employing therapeutic gene sequences encoding t-PA, urokinase, etc.) is delivered via catheter directly upstream of the thrombus-prone segment, or directly to the segment itself (possibly in conjunction with tacking-down a coronary dissection).

4. Adenoviral Delivery to Cardiac Muscle Cells

Since the ventricular myocytes do not divide, retroviral vectors (which require host cell proliferation) cannot be employed. Adenoviruses, however, infect non-replicating cells, and exhibit a significant ability to mediate more extensive gene delivery in skeletal muscle than plasmid DNA. Thus, they are suitable for use in cardiac muscle gene therapy by direct or intravenous delivery. Moreover, recombinant adenovirus mediates remarkably uniform gene transfer to adult cardiac cells in culture. Adenovirus-mediated gene transfer to the adult ventricular myocardium has been demonstrated in vivo, by local injection or coronary artery infusion (Leinwand, L.A. et al., J. Cell. Biochem. 17D:193 (1993); Barr, E. et al., J. Cell. Biochem. 17D:195 (1993), both herein incorporated by reference).

Gene therapies to the myocardium include (but are not limited to) antioxidants (such as superoxide dismutase, catalase, and glutathione peroxidase) to protect against ischemia or reperfusion injury, angiogenic growth factors (such as acidic and basic FGF) to stimulate angiogenesis to ischemic zones of myocardium, growth factors to stimulate cellular proliferation in infarcted areas, and rejection

-25-

inhibitors (such as CD8) to prevent immune rejection of transplanted hearts.

5. Adenovirus-mediated Gene Therapy for Inflammation

5 For the treatment of inflammatory disorders (such as reperfusion injury of myocardial or other tissues, etc.) suitable therapeutic gene sequences include those that encode antisense molecules or therapeutic proteins that can suppress the expression of ICAMs such as ICAM-1
10 or ICAM-2 (Rothlein *et al.*, J. Immunol. 137:1270 (1986); European Patent Application Publication No. 289,949; Staunton, D.E. *et al.*, Cell 52:925-933 (1988); Staunton, D.M. *et al.*, FASEB J. 3:a446 (1989); which references are incorporated herein by reference), VCAM-1 (Hynes,
15 R.O. Cell 48:549-554 (1987)) or selectins (such as L-selectin (also termed LECCAM-1, MEL-14, LAM-1, LECAM-1 or lymphocyte homing receptor); E-selectin (also known as endothelial leukocyte adhesion molecule-1 (ELAM-1"), or LECCAM-2); and P-selectin (also known as CD62,
20 platelet activation dependent granule external membrane (PADGEM), LECCAM-3, or granule membrane protein-140 (GMP-140); see, Gallatin, M.W. *et al.*, Nature 304:30-34 (1989); Lasky, L.A. *et al.*, Cell 56:1045-1055 (1989); Siegelman, M.H. *et al.*, Proc. Natl. Acad. Sci. (U.S.A.)
25 86:5562-5566 (1989); Tedder, T.F. *et al.*, J. Exper. Med. 170:123-133 (1989); Bevilacqua, M.P. *et al.*, Science 243:1160-1165 (1989); Bevilacqua, M.P. *et al.*, Proc. Natl. Acad. Sci. (U.S.A.) 84:9238-9242 (1987); Iuscinskas, F.W. *et al.*, J. Immunol. 143:3318-3324
30 (1989); Pober, J.S. *et al.*, Lab. Invest. 64:301-305 (1991)).

B. The Preferred Vectors of the Present Invention

As used herein, the term "vector" is intended to denote a nucleic acid molecule. Preferably, such vectors will include therapeutic gene sequences and viral or plasmid DNA. A gene transfer vector is a virus or synthetic complex that contains a therapeutic gene sequence, and is able to mediate the transfer of the gene sequence into a recipient cell. Most preferably, the vectors of the present invention are associated with, or contained in a carrier, which may be a liposomal complex, a viral particle, etc. In such a case, the vectors are referred to as liposomal vectors, or viral vectors.

Adenoviruses, especially modified adenoviruses, are the preferred viral vector for delivering the therapeutic gene sequences of the present invention. Adenoviruses are intermediate-sized DNA viruses with genomes consisting of linear double-stranded DNA molecules of approximately 36,000 bp. The virion is an icosahedron about 70 nm in diameter, consisting exclusively of protein and DNA. Adenoviruses have been isolated from a large number of different species (mammalian and fowl) and over 100 different serotypes have been reported, some 43 of them human. The human adenoviruses, particularly types 2, 5, and 12, have been the most extensively characterized, and these viruses have served as valuable tools in the study of the molecular biology of DNA replication, transcription, RNA processing, and protein synthesis in mammalian cells (for review, see, Ginsberg, H., "The Adenoviruses," Plenum Press, NY (1984); Berkner, K.L. et al., Biotechniques 6:616-629 (1988)). The biology of adenoviruses is reviewed by Graham, F.L. et al. (In: Methods in Molecular Biology: Gene Transfer and Expression Protocols, E.J. Murray, Ed. (Humana Press,

-27-

New Jersey, (1991), Vol. 7, chap. 11, pp. 109-128 herein incorporated by reference).

Adenoviruses have several salient advantages over other gene therapy vectors. The viral particle is
5 relatively stable, and, in the case of serotypes commonly used as vectors to date, the viral genome does not undergo rearrangement at a high rate. Insertions of foreign genes are generally maintained without change through successive rounds of viral replication. The
10 adenovirus genome is also relatively easy to manipulate by recombinant DNA techniques, and the virus replicates efficiently in permissive 293 host cells.

Unlike retroviral vectors, adenoviral vectors do not require host cell replication in order to achieve
15 high-level expression. This is due to the fact that adenoviral DNA remains episomal and does not need to integrate into the host genome before becoming transcriptionally active. Thus, adenovirus extends the range of cells and tissues that might be amenable to genetic manipulation, including the principal potential
20 targets for cardiovascular applications.

The key advantage to the use of adenoviral vectors is the dramatic efficiency with which they mediate gene transfer. This efficiency is primarily due to the
25 molecular mechanisms by which adenovirus mediates membrane penetration, lysosomal escape, and nuclear entry (Marsh, M. et al., In: Advances in Virus Research, K. Maramorosch et al., Eds. (Academic Press, New York, 1989), vol. 36, p. 107-151, herein incorporated by
30 reference). In contrast, Lipofectin™-DNA complexes are only efficient at achieving membrane fusion; the vast majority of the DNA is then degraded by the lysosomes, and very little DNA ever reaches the cellular nucleus. Adenoviral vectors carrying antiproliferative genes can
35 be directly targeted to a defined site, such as a site of angioplasty using a perforated infusion catheter, thereby ensuring maximal gene expression at the precise

-28-

site where it is required. Adenoviruses are equally effective in mediating gene transfer into human, porcine, and canine smooth muscle cells, and the kinetics of viral binding are such that even a brief exposure to adenovirus will result in significant levels of gene transfer.

Most preferably, a modified adenoviral vector will be used to deliver the therapeutic gene sequences. The most preferred adenoviral vector is one that is replication-deficient due to deletion of critical genes in the E1 region of the viral genome. Such vectors can only be propagated in cell lines such as the permissive 293 host cell line, which provides the necessary E1a and E1b gene products in trans (Graham, F.L. et al. (In: Methods in Molecular Biology: Gene Transfer and Expression Protocols, E.J. Murray, Ed. (Humana Press, New Jersey, (1991), Vol. 7, chap. 11, pp. 109-128). Whereas wild-type Ad5 (containing E1 genes, and competent for viral replication in cells) can be cytopathic within 6 to 48 hours, the deletion of the E1 genes in the replication-deficient Ad5 vectors of the present invention precludes cytopathy.

The absence of chromosomal integration means that the DNA from a replication-deficient adenovirus has a finite lifespan (several weeks or more) before it is ultimately degraded by host nucleases. Thus, it is ideally designed for the treatment of cardiovascular disorders such as restenosis, and other acute diseases and conditions. As shown below, the time course of recombinant gene expression from such adenoviral vectors matches the time frame necessary to deliver antiproliferative therapy in vivo. The gradual decline in expression which is observed after two weeks is actually advantageous, since the expression of antiproliferative proteins at later time points might be inappropriate for maintenance and integrity of the arterial wall.

-29-

The replication-deficient adenoviral vectors currently utilized for in vivo gene transfer are largely derived from adenovirus serotype 5 (Ad5), although a few contain elements from closely related serotypes. Wild-type Ad5 is an icosahedral, non-enveloped virus carrying a linear, double-stranded 36 kilobase DNA genome. The molecular genetics and virology of Ad5 have been well characterized, so that much is known about Ad5 genes, their expression, and their function in viral replication (Ginsberg, H., "The Adenoviruses," Plenum Press, NY (1984)). For the cardiologist, the critical properties of Ad5 are those which favor its use for gene transfer in vivo -- the availability of replication-deficient derivatives, the comparative ease with which the Ad5 genome can be manipulated, the variety of methods for virus delivery, the efficiency for infecting mammalian cells, the high levels of recombinant protein expression which can be obtained, and the relative lack of cytotoxicity. Adenovirus can be concentrated to exceptionally high titers, exceeding 10^{10} pfu/ml, can be engineered to accommodate very large DNA inserts, and possesses a broad host range, enabling tests in diverse animal models of human disease. Finally, the DNA copy generated from a retrovirus's RNA genome becomes permanently and randomly integrated into some chromosomal site within infected cells, a process that can give rise to aberrant cell behavior. The adenoviral genome, by contrast, remains physically separate from host cell DNA (episomal), obviating this hazard of insertional mutagenesis. As additional merits regarding safety, human adenoviruses have not been associated with clinical malignancies and, indeed, live adenovirus vaccines have been administered to large numbers of subjects without ill effect. Replication-deficient adenoviral vectors have been used to mediate in vivo gene transfer into bronchial epithelium (Rosenfeld, M.A. et al., Cell 68:143-155 (1992)) and skeletal muscle

-30-

(Quantin, B. et al., Proc. Natl. Acad. Sci. (U.S.A.)
89:2581-2584 (1992)).

5 In a particularly preferred embodiment, the
adenoviral vectors will be further modified so as to
carry ligands specific for receptors present on the
desired target cells. Alternatively, synthetic vectors
can be constructed from synthetic and/or recombinant
products. The synthetic vectors would be constructed
of: i) one or more recombinant DNA cassette(s) directing
10 the expression of therapeutic gene sequence(s)
(preferably, but not necessarily in the form of
plasmid(s)), ii) a ligand to mediate cell attachment
and/or endocytosis (preferable a ligand which targets a
specific cell or tissue), iii) a component for
15 mediating escape from the lysosomal pathway (such as the
hemagglutinin spike protein from the influenza virus),
and iv) a nuclear localization signal (NLS) peptide or
other mechanism for mediating transport to the nucleus.
The high efficiency gene transfer provided by these
20 methods is described in Example 8.

These components can be linked together by
covalently attaching polycationic linkers (such as poly-
L-lysine) to the non-DNA components, then mixing these
linked components with the DNA component(s) to form
25 ionic complexes. Additionally or alternatively, labile
covalent bonds can be utilized which will break under
specific conditions (such as under low pH in lysosomes).
Additionally or alternatively, liposomal complexes of
cationic lipids (i.e. liposomes, or lipid-protein
30 complexes) can be used to maintain association of the
DNA-based components with the other functional
components. Additionally, modifications can be made to
the DNA-based component in order to prolong its
expression in the nucleus. Such modifications include
35 the chemical modification of the DNA in order to prevent
its degradation, and/or providing a means of DNA
replication, such as including the Epstein-Barr virus

-31-

(EBV) origin of replication (ori-P) and the gene encoding EBV nuclear antigen 1 (EBNA-1)).

The capacity and efficiency of the vectors of the present invention to mediate gene therapy can be effectively demonstrated by incorporating a marker gene in place of the therapeutic gene sequence, and then determining the kinetics of marker gene expression in the recipient patient. The luciferase reporter system is a preferred marker gene. Alternatively, since each reporter system has its own unique and useful properties, other marker genes may be employed. One feature of note in the luciferase reporter system is the short half-life of the protein in the cell. Thus luciferase does not accumulate to high levels in vitro or in vivo due to its rapid turnover in cells. In contrast, the β -galactosidase gene product of the *E. coli* LacZ reporter gene is remarkably stable in eucaryotic cells. The histochemical X-gal stain for lacZ positive cells is compromised by cells within the arterial wall which express the mammalian β -galactosidase activity. This problem can be overcome by adding sequences encoding a nuclear localization signal (NLS) to the 5'-end of the *E. coli* LacZ gene. After transcription and translation of this fusion gene, the NLS directs transport of β -galactosidase to the nucleus. Following fixation and incubation with the X-gal substrate, the distinct nuclear staining will definitively identify the cells which carry the reporter gene. As predicted by their relative stabilities, we have found that β -galactosidase accumulates to higher levels than luciferase after gene transfer, both in vitro and in vivo. Thus the two reporter systems complement each other by indicating the range of gene expression that might be expected from a therapeutic construction after gene transfer. The absolute levels of therapeutic gene expression obtained in vivo will depend upon a large number of variables including: DNA

-32-

copy number, DNA stability, promoter strength, transcription rate, messenger RNA stability, translation rate, post-transnational modifications, and protein stability.

5 **III. The Administration of the Molecules of the Present Invention**

 The genetic therapy of the present invention can be administered in a variety of ways. In one embodiment, such therapy can be provided alone, i.e as the sole
10 therapy administered to a patient. Alternatively, two or more different therapeutic gene sequences can be provided to a patient in a manner that results in overlapping expression of more than one introduced therapeutic gene sequence. For example, a patient can
15 be administered a combination of therapeutic vectors that encode two or more of the following therapeutic gene sequences: Thrombin inhibitors (factor Xa antagonists, hirudin, thrombomodulin, antithrombin III, protein C, protein S, lipoprotein-associated coagulation
20 inhibitor), Platelet inhibitors (RGD peptides, GP IIb/IIIa receptor antagonists), GF inhibitors (calcitonin gene-related peptide, somatostatin, soluble growth factor receptors, dominant-negative receptors, receptor antagonists), Antioxidants (superoxide
25 dismutase, catalase, glutathione peroxidase), Antiproliferative proteins (tyrosine phosphatase, γ -interferon, p53, elF-2 kinase), Chimeric toxins (FGF-saporin, EGF-diphtheria toxin, TGF α -pseudomonas exotoxin A), Anti-inflammatory agents, TGF β , lipocortin,
30 vasocortin, IL-10), Antilipemic agents (LDL receptor, ApoA1, phospholipases) and Antisense or Ribozyme RNAs targeted against: c-myc, c-myb, c-fos, c-jun, PCNA, tyrosine kinase receptors, nucleolar p120 protein, slow calcium channels, HMG-CoA reductase, cholesterol
35 acyltransferase, cholesterol ester transfer proteins,

-33-

thromboxane synthase, thromboxane receptors, etc. The combination of two or more different classes of therapeutic sequence, such as hirudin (a thrombin inhibitor) and γ -interferon (an antiproliferative), may effect a more potent or synergistic therapy for restenosis. Similarly, a gene therapy comprising sets of vectors that mediate the expression of a solubilized ICAM-1 and a solubilized selectin can be employed. In such combined vector therapies, the multiple therapeutic gene sequences can be present on the same introduced vector, or can be present on separate vectors since a combination of vectors can be delivered as easily as a single vector.

As indicated, the time course of the genetic therapy of the present invention commences within hours of administration, and lasts for several weeks or longer. In a preferred embodiment, such gene therapy can be used in combination with conventional pharmacological therapies to effect a more potent or synergistic response. Thus, conventional anti-inflammatory agents, anti-restenotic agents, anti-thrombin factors, etc. can be used in combination with one or more of the therapeutic gene sequences of the present invention. The utility of combining such therapies lies in the complementarity in the intervals over which each of the therapies are effective. After administration, a pharmaceutical is generally effective for a period of hours, and is seldom effective for more than one day. Conversely, the genetic therapy of the present invention may take hours to become effective, and will continue to be effective over a period of time far in excess of that which can be obtained by pharmaceutical therapy.

A. In Vivo Administration

Any of a variety of methods can be used to provide the therapeutic vector molecules to a desired site. A desirable coronary delivery system has the general properties of being atraumatic, and permitting a prolonged contact of relatively small volumes with the arterial wall, while maintaining adequate distal perfusion.

In one embodiment, the therapeutic gene sequences are provided to a localized, preselected site using percutaneous catheter-based local delivery systems. One such device (the Wolinsky Infusion Catheter made by USCI) insures gene delivery to the vessel wall, but occasionally causes intramural hematomas and dissections that are grossly evident at three days post infusion. The severity of injury can be reduced by choosing appropriately sized balloons (balloon to artery ratio of 1:1) and lower infusion pressures (2-3 atmospheres). However, the kinetics of viral binding and the high efficiency of gene transfer afforded by adenoviral vectors decreases any need for high pressure delivery, and enables adequate levels of gene expression to be achieved with sustained, low pressure infusions which should not impose significant arterial damage.

In lieu of using such catheters, other catheter designs may be employed including: i) microporous balloons (made of dialysis-like membranes of defined porosity which allow the infusate to seep out rather than jet out), ii) hydrogel-coated balloons (in which the therapeutic component is incorporated into the hydrogel for distribution into the arterial wall), iii) dual balloons (with infusion port between the two balloons), iv) double-balloons (with one balloon inside a second perforated or microporous balloon), v) channel balloons (in which the distribution of the therapeutic agent is compartmentalized), and vi) seeded balloons (in

-35-

which inflation causes the rupture of "seeds" which release therapeutic reagents into the vascular wall).

Gene expression can preferentially be targeted to the luminal surface of the artery by low-pressure, dual-balloon delivery; or to the deeper layers in the artery by high-pressure, Wolinsky balloon delivery. Thus the catheter delivery device, as well as the manner in which it is employed, can be used to select and determine the cellular populations which are most likely to receive the recombinant genes.

In an alternative embodiment, the therapeutic gene sequences are provided to a localized, preselected site using an implanted stent, graft or implant, that is introduced into the desired site, as by interventional or surgical procedures, etc. Control release preparations may be achieved through the use of polymers to complex or absorb the adenoviral vectors. The controlled delivery may be exercised by selecting appropriate macromolecules (for example polyesters, polyamino acids, polyvinyl, pyrrolidone, ethylenevinylacetate, methylcellulose, carboxymethylcellulose, or protamine sulfate) and the concentration of macromolecules as well as the methods of incorporation in order to control release. Another possible method to control the duration of action by controlled release preparations is to incorporate the adenoviral vectors into particles of a polymeric material such as polyesters, polyamino acids, hydrogels, poly(lactic acid) or ethylene vinylacetate copolymers. Alternatively, instead of incorporating these vectors into polymeric particles, it is possible to entrap these materials in, for example, hydroxymethylcellulose, gelatin, etc. Such techniques are disclosed in Remington's Pharmaceutical Sciences (1980).

B. Ex vivo Administration

In an alternative embodiment, the vectors of the present invention may be provided to patients using an ex vivo regimen. This ex-vivo approach to targeting gene therapy to a particular organ, tissue, and/or cell population has a unique advantage since the organ/tissue/cell can be subjected for extended periods of time to high concentrations of gene transfer vector resulting in high-efficiency gene transfer. For example, vascular grafts such as saphonous veins or mammary arteries can be treated with the gene transfer vectors elaborated herein between the explantation and implantation procedures to prevent the accelerated atherosclerosis and restenosis which confounds autologous vascular graft transplantations such as coronary artery bypass grafts. A second application is in the field of cardiac transplantation, where the explanted heart could be bathed in a cardioplegic solution containing a gene transfer vector which prevents transplant rejection (such as by expressing CD8 to inhibit allospecific T-cell responders, etc.) or which mediates against ischemic and/or reperfusion injury (by directing the expression of antioxidants including superoxide dismutase, catalase, glutathione peroxidase, etc.). A third ex-vivo example involves liver transplantation. In this case, an explanted liver destined for transplantation into a hyperlipidemic person could be modified with the gene transfer vectors described herein to produce additional LDL receptors and/or apolipoprotein A1, etc. The examples detailed above are utilized for illustrative purposes and should in no way limit the general approach of genetically modifying explanted organs/tissues/cells before implantation into the patient. In some cases, it may even be desirable to explant organs/tissues/cells solely for the purpose of obtaining high-efficiency gene

-37-

transfer before replacing them back into the same patient.

Having now generally described the invention, the same will be more readily understood through reference to the following examples which are provided by way of illustration, and are not intended to be limiting of the present invention, unless specified.

Example 1

Direct In Vivo Gene Transfer: Influences of Delivery Volume and Pressure

In order to evaluate the basic parameters which influence the efficiency of direct gene transfer, a perforated balloon delivery system (Wolinsky Infusion Catheter [WIC], USCI, Billerica, MA) was employed. 50 μ g of reporter vector DNA (pRSVL) was complexed with 150 μ g of Lipofectin™ (BRL, Gaithersburg, MD) and diluted to either 4 or 8 ml with Opti-MEM I® (also from BRL). The volumes were infused into the RCA, LAD, and LCX coronary arteries of 8 intubated Hanford miniature swine using modestly oversized WIC's (balloon/artery ratio of 1.1-1.3) and inflation pressures of either 4 or 8 atm. The animals were revived following catheterization and maintained for 3 days before sacrifice for arterial harvest and luciferase assay using a Monolight 2010 luminometer (Analytical Luminescence Laboratory [ALL], San diego, CA). Background was subtracted and the total mass of luciferase protein recovered from each artery was calculated after calibration of the luminometer using certified luciferase standards (ALL). Luciferase activity corresponding to ≥ 3 fg protein was detected in 23 of 24 arteries (96%) with a mean of 283 fg luciferase recovered per infused artery. The results of varying the DNA concentrations and balloon pressures are shown in Table 1, where each value in the 2 x 2 matrix gives the mean femtograms (fg) of active luciferase recovered

-38-

from 6 arteries (2 RCAs, 2 LADS, 2 LCXS) \pm standard deviation.

TABLE 1 EFFECT OF VOLUME AND PRESSURE		
	4 atm	8 atm
8 ml	43 \pm 44	148 \pm 155
4 ml	216 \pm 368	771 \pm 937

Thus, using equal amounts of reporter vector, both the higher DNA concentration (4 vs 8 ml delivery volume), and the higher infusion pressure (8 vs 4 atmospheres) resulted in more efficient gene transfer, with maximum transfer achieved when these two parameters were combined.

Example 2

Construction of an Adenoviral Reporter Vector Carrying the Luciferase Gene

The lacZ reporter gene has been employed in direct gene transfer experiments targeting peripheral (Nabel, E.G. et al., Science 249:1285 (1990); Flugelman, M.Y. et al., Circulation 85:1110-1117 (1992); Lemarchand, P. et al., Circ. Res. 72:1132-1138 (1993)) and coronary arteries (Lim, C.S. et al., Circulation 83:2007-2011 (1991)), but endogenous β -galactosidase activity has been observed in arterial sections by our laboratory and others (Lim, C.S. et al., Circulation 83:2007-2011 (1991); Flugelman, M.Y. et al., Circulation 85:1110 (1992)). For this reason, the luciferase reporter system was used to accurately quantitate levels of recombinant protein produced in coronary arteries following direct gene transfer.

Early experiments employed complexes of Lipofectin™ (a 1:1 (w/w) liposome formulation of a cationic lipid and dioleoyl phosphatidylethanolamine obtained from BRL;

-39-

Gaithersburg, Maryland) and a reporter plasmid (pRSVL) which carries the long terminal repeat of the Rous sarcoma virus, the firefly luciferase cDNA, and the SV40 small-t intron / polyadenylation signal (Keller, G.A. et al., Proc. Natl. Acad. Sci. (U.S.A.) 84:3264-3268 (1987)). The inclusion of 1 mg/ml bovine serum albumin (BSA) in the Lipofectin™ cocktail results in higher and more reproducible results, presumably because it acts as a carrier and thus prevents the DNA-Lipofectin™ complexes from adhering to the plastic tubes, pipettes and catheters. The addition of influenza hemagglutinin protein (HA) yields results that are similar to those obtained with BSA. However, combining HA with BSA in the transection cocktail failed to yield further improvements in gene transfer. Because sub-physiological levels of recombinant gene expression were obtained using Lipofectin™, a replication-deficient adenoviral vector carrying an analogous expression cassette was constructed (Figure 1).

As shown in Figure 1, the adenoviral vector carrying the luciferase gene was derived using the replication-deficient adenovirus type 5 (Ad5) cloning system of F.L. Graham (McGrory, W.J. et al., Virol. 163:614-617 (1988), herein incorporated by reference). Briefly, a firefly luciferase expression cassette analogous to the one in pRSVL (Keller, G.A. et al., Proc. Natl. Acad. Sci. (U.S.A.) 84:3264-3268 (1987), herein incorporated by reference) was subcloned between left-end adenoviral sequences in pXCJL.1 to create pXCJL.1/RSV/GL2. This construction was cotransfected into the transformed 293 embryonic kidney cell line (Graham, F.L. et al., J. Gen. Virol. 36:59-72 (1977)) with a plasmid (pJM17) which carries a circularized dl309 adenoviral genome (McGrory, W.J. et al., Virol. 163:614-617 (1988)), herein incorporated by reference). The presence of both the PBR322 derivative (pBRX) and adenoviral genome within pJM17 causes this plasmid to

Putative viral clones were plaque purified, screened for the presence of the luciferase gene insert, propagated, isolated and titered according to the protocols of Graham, F.L. et al. (In: Methods in Molecular Biology: Gene Transfer and Expression Protocols, E.J. Murray, Ed. (Humana Press, New Jersey, (1991), Vol. 7, chap. 11, pp. 109-128).

20 Example 3
Comparative Efficiency of
Lipofectin™- vs. Adenovirus-mediated Gene Transfer

To further characterize the comparative efficiency of replication-deficient adenoviral vectors vs Lipofectin™ at mediating direct gene transfer into living coronary arteries, 50 µg of reporter vector DNA (pCMVβ, in which the CMV promoter drove the expression of the E. coli LacZ gene; or pRSVL, in which the long terminal repeat (LTR) from the Rous sarcoma virus (RSV) drove the expression of the firefly luciferase cDNA) was complexed with 150 µg of Lipofectin™ (BRL) and diluted to 4 ml with Opti-MEM I (BRL).

Viral infections were performed with 4 ml dilutions of recombinant adenoviruses harboring analogous expression cassettes: 5 x 10⁹ pfu per artery of

-41-

Ad5/HCMV/LacZ (in which LacZ is expressed from the CMV promoter; vector obtained from A. Bett and F.L. Graham) or 4×10^9 pfu per artery of Ad5/RSV/GL2 (in which the luciferase gene is expressed from the RSV LTR). Under fluoroscopy, these solutions were infused at a pressure of 8 atmospheres into the coronary arteries of intubated Hanford miniature swine using perforated balloon catheters. Levels of reporter gene activity were determined 3 days following infusion catheterization using a chemiluminescent assay for β -galactosidase and a luminescent assay for luciferase. The results of this comparison are shown in Table 2, where each value in the 2x2 matrix represents the mean amount of reporter gene product recovered from 3 to 6 coronary arteries \pm standard deviation.

TABLE 2 COMPARATIVE EXPRESSION EFFICIENCY		
Reporter Protein	β -galactosidase (ng)	Luciferase (pg)
Lipofectin™	1.52 ± 0.68	0.11 ± 0.10
Adenovirus	106 ± 96	35.9 ± 27.7

Thus, with both reporter systems, markedly higher levels of recombinant gene product were obtained from adenoviral vectors than from Lipofections using 50 μ g of highly purified plasmid DNA. The absolute amounts of recombinant gene product obtained were obviously dependent upon the mass of DNA (or titer of virus) employed; nevertheless, this study demonstrated that when equal volumes were infused, replication-deficient adenoviral vectors were far more efficient than Lipofectin™ at mediating direct gene transfer into living coronary arteries. A 70-fold difference was observed using the LacZ reporter gene, and a 240-fold difference was observed using luciferase. It is

-42-

interesting to note that the nanogram levels of β -galactosidase were far in excess of the picogram levels of luciferase. Most of this difference is probably due to the high stability of the β -galactosidase protein relative to luciferase, although promoter strength may also be a contributing factor. Nevertheless, both reporter systems demonstrate the clear superiority of adenoviral vectors over more conventional methods of gene transfer.

10

Example 4Direct Gene Transfer by Adenoviral Vectors:
Time Course of Therapeutic Gene Sequence Expression

Multiple lines of evidence, from animal models as well as human autopsies, indicate that the majority of the smooth muscle cell migration and proliferation responsible for restenosis occur during the first two weeks following coronary angioplasty. Thus in order to be most effective, an antiproliferative gene therapy directed against restenosis would have to provide for at least two weeks of recombinant gene expression. It was therefore important to determine if the duration of recombinant gene expression provided by these vectors makes them suitable for use in a genetic therapy, such as one against restenosis. Under fluoroscopy, a 4 ml volume containing 4×10^9 pfu of Ad5/RSV/GL2 was infused into each porcine coronary artery at a pressure of 8 atmospheres using a Wolinsky catheter. Luciferase activity in the arteries was determined upon sacrifice at 1, 3, 7, and 14 days following catheterization. The results are reported in Table 3, where each value represents the mean picograms (pg) of active luciferase recovered from 3 to 6 coronary arteries \pm standard deviation (sd).

-43-

TABLE 3 TIME COURSE OF EXPRESSION				
	Time From Infusion to Assay (Days)			
	1	3	7	14
pg luciferase /artery ± 1 sd	24±23	27±21	21±39	12±19
No. of Coronaries Assayed	3	6	6	6

5 The experiment indicates that similar levels of gene expression are obtained at 1, 3, 7, and 14 days following gene transfer, with a trend towards maximal
10 expression at 3-days. Significantly, the results also indicate that gene expression could be detected after only a single day, indicating that gene therapy will have a therapeutic effect within 24 hours. Expression has been observed even at 28 days following gene
15 transfer. This study provides strong evidence to suggest that the duration of gene expression provided by adenoviral vectors is suitable for antiproliferative gene therapies directed against coronary restenosis.

20 The luciferase assays three days after gene transfer further demonstrated that 4×10^9 plaque-forming units of recombinant virus produced at least 100-fold more luciferase than 50 μ g of the analogous plasmid DNA-Lipofectin™ complex (Mazur, W. *et al.*, *J. Cell. Biochem.* 17E:226 (1993), herein incorporated by reference).
25 Comparable levels of recombinant gene expression were obtained in vivo as early as 24 hours after gene transfer, and expression persisted for at least 14 days (Ali, M.N. *et al.*, *Restenosis Summit V*, May 20-21, Cleveland, OH (1993), herein incorporated by reference).
30 Thus, both the magnitude and duration of gene expression provided by recombinant adenoviruses appear suitable for the prevention of coronary restenosis.

-44-

Example 5Direct Gene Transfer by Adenoviral Vectors:
Efficiency of Expression in the Major Coronary Arteries

A comparison was made of Lipofectin™- and
5 adenovirus-mediated gene transfer in all three major
porcine coronary arteries: the circumflex (LCX), left
anterior descending (LAD), and right coronary arteries
(RCA). Male Hanford miniature swine weighing 50-100
10 lbs. at exsanguination were employed in a protocol which
adhered to NIH and institutional guidelines.

Direct gene transfer into normal porcine coronary
arteries was carried out under fluoroscopic guidance
using a strictly percutaneous approach. All three major
coronary arteries in each animal were subjected to gene
15 transfer using Wolinsky Infusion Catheters (USCI;
Billerica, MA) oversized by 1.1-1.3 times the diameter
of the target arterial segment. These catheters
resemble conventional angioplasty devices except that
the balloon is perforated with 28 microscopic holes
20 (each 25 μ m in diameter). The infusion balloon was
positioned at the target site in each coronary, 4 ml of
gene transfer reagent was loaded into the inflation
device, and a pressure of 8 atmospheres was applied;
resulting in simultaneous balloon inflation and gene
25 infusion over a period of 12 to 28 seconds. For
Lipofectin™ transfection of each coronary, 50 μ g of
pRSVL (Keller, G.A. et al., Proc. Natl. Acad. Sci.
(U.S.A.) 84:3264-3268 (1987)) was complexed with 150 μ g
of Lipofectin™ and diluted to a final volume of 4 ml
30 with OptiMEM-I® (BRL). For each adenoviral infection, 4
 $\times 10^9$ plaque forming units of Ad5/RSV/GL2 (Figure 1),
were diluted to 4 ml with OptiMEM-I®.

Following gene transfer and angiography, the
procedure was concluded and the swine were maintained
35 for three days before being placed under deep anesthesia
for exsanguination and analysis for luciferase
expression. For such procedures, hearts were removed,

-45-

placed in ice-cold PBS, and coronary arteries dissected from myocardium. The vessels were divided into multiple segments (each <1 cm long), minced, and homogenized in 100 μ l of luciferase lysis reagent (25 mM Tris-phosphate, pH 7.8. 2 mM DTT, 2 mM 1,2-diaminocyclohexane-N,N,N',N'-tetraacetic acid, 10% glycerol, and 1% Triton X-100). The samples were centrifuged at 12,000 g for 5 minutes, and duplicate luciferase assays were performed with a Monolight 2010 luminometer (Analytical Luminescence Laboratory [ALL]; San Diego, CA). 100 μ l of assay reagent (20 mM Tricine, pH 7.8, 1.07 mM (MgCO₃)₄Mg(OH)₂·5H₂O, 2.67 mM MgSO₄, 0.1 mM EDTA, 33.3 mM DTT, 270 μ M coenzyme A, 470 μ M luciferin, and 530 μ M ATP) was added to each 20 μ l sample of cleared supernatant and photons were counted for 30 seconds. Background values from vehicle-infused arteries (equivalent to background from lysis reagent) were subtracted and relative light units were converted to mass units (pg) using calibration curves generated from parallel reactions performed with certified luciferase control standard (ALL).

Using these techniques, the efficiency of Lipofectin™- vs. adenovirus-mediated in vivo gene transfer was evaluated with respect to the type of coronary artery undergoing gene transfer. The total amounts of luciferase recovered from each vessel undergoing the two types of gene transfer are reported in picograms (pg) in Table 4 beneath the type of artery undergoing gene transfer: RCA, right coronary artery; LAD, left anterior descending; LCX, circumflex.

-46-

TABLE 4 NORMAL CORONARY ARTERIES			
Animal #	RCA	LAD	LCX
<u>Transfected by Lipofectin™ (pg)</u>			
39	0.08	0.16	0.02
43	0.28	0.06	0.06
<u>Infected by Adenovirus (pg)</u>			
31	67.9	19.5	20.4
51	10.9	13.5	27.8

Variability in the levels of gene expression was observed using either Lipofectin™- or adenovirus-mediated gene transfer. In normal porcine coronary arteries, luciferase levels obtained with Lipofectin™ ranged from 0.02-0.28 pg, while levels obtained with the adenovirus ranged from 11-68 pg. Thus, the mean level of gene expression with adenovirus in the normal coronary arteries was 240 times higher than that achieved with Lipofectin™. The number of arteries of a certain type undergoing a particular gene transfer regimen is small (n=2), and the results are variable; nevertheless, it is also evident that adenovirus-mediated gene transfer yields similar results, regardless of the type of coronary artery targeted (RCA, LAD, or LCX).

The study summarized above was designed to compare the efficiency of Lipofectin™- versus adenovirus-mediated gene transfer when relatively small volumes (4 ml) containing average concentrations of DNA (or virus) were infused under identical conditions. The absolute levels of luciferase obtained from either gene transfer method can be improved by increasing the concentration of reporter gene or the volume of infusate. For example, Chapman, G.D. *et al.* (*Circ. Res.* 71:27-33 (1992)) achieved picogram levels of luciferase in canine coronary arteries using Lipofectin™ and similar percutaneous catheters, but the amounts of plasmid DNA

-47-

(up to 650 μ g) and the volumes of infusate (up to 20 ml) were far greater than those used here. Analogous strategies for adenoviral vectors (such as increased titer) should result in levels of luciferase production far higher than those reported here. Of importance to the long-term clinical application of such methods, adenovirus-mediated gene transfer was found to be comparably effective in normal and atherosclerotic arteries, and vessels subjected to balloon injury or stent deployment.

Example 6

Direct Gene Transfer into Atherosclerotic Coronary Arteries

The results in Table 4 demonstrate efficient gene transfer by the adenoviral vector into physiologically normal coronary arteries. Since the vascular targets for gene therapy against restenosis include atherosclerotic arteries and those displaying the pathology of VSMC hyperplasia, the efficiency of gene transfer was examined in three types of atherosclerotic porcine coronary arteries: uninjured, injured by oversized balloons, and injured by intracoronary stents (Table 5).

Hypercholesterolemia was induced with a high fat diet which produces predictable atherosclerotic changes in the coronary arteries (Pond, W.G., In: Swine in Cardiovascular Research, H.C. Stanton and H. J. Morsmann, Eds. (CRC Press, Florida, 1986), Vol, 2. chap. 1, herein incorporated by reference). The additional manipulations involving injury by oversized balloons (Schwartz, R.S. et al., Circulation 82:2190 (1990) herein incorporated by reference) or intracoronary stents (Rodgers, G.P. et al., Circulation 82:560 (1990); Karas, S.P. et al., J. Amer. Col. Cardiol. 20:467 (1992), both herein incorporated by reference) have been

shown to produce arterial lesions with histopathologies which closely resemble human restenosis.

The atherosclerotic cohort (8 animals total) was initiated on a diet of 2% cholesterol, 15% fat, and 1.5% sodium cholate (Pond, W.G., In: Swine in Cardiovascular Research, H.C. Stanton and H. J. Morsmann, Eds. (CRC Press, Florida, 1986), Vol, 2. chap. 1) one month before catheterization. This cohort had a mean cholesterol level of 282 mg/dl during the first procedure and a mean of 266 mg/dl at sacrifice two months later.

After one month of atherosclerotic diet, the animals were sedated (IM ketamine, 20-30 mg/kg; acepromazine, 0.22 mg/kg; and atropine, 0.05 mg/kg, intubated, and anesthetized with halothane 2%. A femoral artery was isolated and a coronary guiding catheter was inserted then directed to the target coronary artery. In one major coronary, a balloon expandable flexible tantalum stent (oversized by 1.1-1.3 times the vessel diameter) was deployed under angiographic guidance. In another major coronary, balloon overstretch injury was imposed with a balloon catheter oversized by 1.1-1.3 times vessel diameter by inflating three times for 60 seconds at 8-10 atmospheres. During each instrumentation, angiography was performed to insure future identification of the target segment. The third major coronary was not instrumented and served as an uninjured atherosclerotic control. Following the procedure, the animals were maintained on atherogenic diet for two additional months until gene transfer.

The same interventional procedures described above were used to prepare for gene transfer using a strictly percutaneous approach. All three major coronary arteries in each animal underwent gene transfer using Wolinsky Infusion Catheters (USCI; Billerica, MA) oversized by 1.1-1.3 times the diameter of the target arterial segment. For Lipofectin™ transfection of each

-49-

coronary, 50 μ g of pRSVL (Keller, G.A. et al., Proc. Natl. Acad. Sci. (U.S.A.) 84:3264-3268 (1987)) was complexed with 150 μ g of Lipofectin™ and diluted to a final volume of 4 ml with OptiMEM-I® (BRL). For each

5 adenoviral infection, 4 x 10⁹ plaque forming units of Ad5/RSV/GL2 (Figure 1), were diluted to 4 ml with OptiMEM-I®. The infusion balloon was positioned at the previously injured target site in each coronary, 4 ml of

10 gene transfer reagent was loaded into the inflation device, and a pressure of 8 atmospheres was applied; resulting in simultaneous balloon inflation and gene infusion over a period of 12 to 28 seconds.

Three days following gene transfer, the swine were placed under deep anesthesia and exsanguinated. The

15 hearts were removed, placed in ice-cold PBS, and coronary arteries dissected from myocardium. The vessels were then assayed for luciferase activity using the same materials and methods described in Example 5 above.

20 Table 5 shows luciferase expression in atherosclerotic porcine coronary arteries including those simulating restenosis. The total amount of luciferase recovered from each atherosclerotic coronary vessel undergoing Lipofectin™- or adenoviral-mediated in

25 vivo gene transfer is reported in picograms (pg) beneath the type of artery undergoing gene transfer: Athero, uninjured atherosclerotic, Ballooned, atherosclerotic arteries injured by oversized balloon; Stented, atherosclerotic arteries injured by intracoronary

30 stents.

TABLE 5 ATHEROSCLEROTIC CORONARY ARTERIES			
Animal #	Athero	<u>Restenotic</u>	
		Ballooned	Stented
<u>Transfected by Lipofectin™ (pg)</u>			
20	0.20	0.47	0.03
08	0.05	0.05	0.02
07	0.39	0.16	0.45
17	0.37	0.14	0.05
<u>Infected by Adenovirus (pg)</u>			
09	5.2	14.9	29.2
15	12.2	3.6	43.4
13	1.4	10.9	26.0
12	8.6	4.3	0.4

Table 5 demonstrates that the high efficiency provided by adenovirus-mediated gene transfer in normal coronary arteries (Table 4), can also be obtained in those with atherosclerotic and restenotic pathology. Figure 2 summarizes the results of direct gene transfer into normal, atherosclerotic and restenotic porcine coronary arteries. In Figure 2, the mean picograms (pg) of active luciferase recovered per coronary artery undergoing each of the eight regimens are indicated on the y-axis. Error bars indicate standard deviation of the mean.

As shown in Figure 2, for either Lipofectin™ or adenovirus, there was no statistically significant difference in the efficiency of gene transfer between the four different types of arteries examined (Normal, Athero, Ballooned and Stented). When all arteries are divided into only two groups (those treated with Lipofectin™ and those treated with adenovirus), the luciferase levels obtained with adenovirus (mean 17.8 pg) were 100 times higher than those obtained with Lipofectin™ (mean 0.169 pg).

In order to obtain a more valid comparison of efficiency, the results were normalized to the number of

-51-

luciferase genes undergoing transfer. Thus 50 μ g of pRSVL DNA (10.6 pmoles of reporter gene) produced a mean 0.169 pg luciferase per coronary artery while 4×10^9 adenoviral particles (6.64 fmoles of reporter gene) produced a mean 17.8 pg luciferase per artery. Therefore, when considered on a molar basis, the adenovirus is nearly 170,000 times more efficient than Lipofectin™ at mediating direct in vivo gene transfer into porcine coronary arteries, and may thus overcome a major obstacle to gene therapy for restenosis. The differential in gene transfer efficiency is due at least in part to the molecular mechanisms by which adenovirus mediates endocytosis, endosomal disruption, and nuclear entry (Marsh, M. et al., In: Advances in Virus Research, K. Maramorosch et al., Eds. (Academic Press, New York, 1989), vol. 36, p. 107-151). In contrast, Lipofectin™-DNA complexes are efficient at penetrating the plasma membrane, but the majority of this DNA is lost, presumably to lysosomal degradation (Ho, S. et al., Cytochem. 31:404 (1983)).

The present results demonstrate that adenoviral vector mediated gene therapy provides far superior results than are obtainable with Lipofectin™. The histochemical study by Nabel, E.G. et al. (Science 249:1285 (1990)) on the direct transfer of the lacZ gene into peripheral vessels suggested that the levels of β -galactosidase obtained with retroviral vectors were comparable to those obtained by Lipofectin™. Similarly, low levels of gene transfer using retroviral vectors were described by Flugelman, M.Y. et al. (Circulation 85:1110 (1992)) in rabbit aortas. Additionally, retroviral vectors may not be well suited for gene therapy against restenosis since: (i) only replicating vascular cells will express the therapeutic gene, (ii) genomic integration by retroviral vectors might interrupt vital cellular genes, or activate inappropriate genes such as cellular oncogenes, and

-52-

(iii) the integrated provirus may continue to express the therapeutic gene for inappropriately long periods of time.

In contrast, replication-deficient Adenovirus 5
5 vectors may be particularly well suited for gene therapy since: (i) they efficiently infect vascular cells regardless of their proliferative state, (ii) the episomal location of the Ad5 genome obviates problems resulting from genomic integration, and (iii) the
10 transient nature of adenoviral infection should result in temporary expression of the therapeutic gene. Permanent expression of therapeutic genes would be unnecessary, and perhaps deleterious, since the threat of restenosis subsides approximately 4 months after
15 angioplasty (Serruys, P.W. et al., Circulation 77:361 (1988)).

In summary, experiments were thus performed with percutaneous catheters comparing the adenovirus with Lipofectin™ in normal arteries, atherosclerotic
20 arteries, and in two atherosclerotic models of restenosis (employing balloon injury or stent deployment). Previous studies have established that direct in vivo gene transfer into the arterial wall was possible (Nabel, E.G. et al., Science 249:1285 (1990)),
25 and that Lipofectin™ and perforated balloon catheters could be used to direct such gene transfer to the coronary arteries of intact animals (Chapman, G.D. et al., Circ. Res. 71:27-33 (1992)). However, the limited efficiency of that technology precluded the possibility
30 of its clinical application. In contrast to such studies, the methods of the present invention demonstrate that these problems can be surmounted in normal, atherosclerotic, and restenotic coronary arteries by using replication-deficient adenoviral
35 vectors to accomplish direct gene transfer in vivo; and thus establishes the feasibility of using these vectors to deliver gene therapy for the prevention of restenosis

-53-

following coronary angioplasty, or to provide gene therapy for other coronary or cardiovascular diseases or conditions.

Example 7

5 Adenovirus-Mediated Direct Gene Transfer
Via Alternative Routes of Administration

10 The examples given above detail percutaneous delivery of gene therapy to coronary arteries via a modified balloon catheter. However, a number of organs/tissues/cells can be targeted by a judicious choice of gene transfer vector and route of administration. For example, luciferase assays of porcine myocardium perfused by coronary arteries undergoing adenovirus-mediated gene transfer demonstrate
15 that the same procedures used to obtain gene transfer into coronary arteries can also be used to obtain gene transfer into defined regions of the myocardium. Reporter gene expression following the direct injection of a gene transfer vector into the myocardium via a
20 hypodermic needle has also been discussed in the scientific literature and confirmed in our laboratory using adenoviral vectors. These routes of administration ensure local delivery of gene transfer reagent for the purposes of confining gene therapy to a
25 particular tissue. However, gene therapies which involve the circulatory system and/or secreted gene products do not necessarily have to be delivered to the coronary arteries or to the myocardium. For example, infusion of adenoviral vectors into the portal vein has
30 been reported to result in high efficiency gene transfer to the liver.

 In order to determine the pattern of gene transfer resulting from direct intraperitoneal or intra-arterial injection of recombinant adenovirus, the carotid
35 arteries or the peritoneal cavities of 1-2 day-old rats

-54-

were directly injected with adenoviral vectors via 27 gauge needles. The rats were sacrificed two days later under deep anesthesia, and seven tissue types were homogenized for luciferase determination. The relative light units of luciferase activity were normalized to the mass of tissue in the assay, and the results are presented in Figure 3. The lower graph of Figure 3 demonstrates that the intraperitoneal injection of Ad5/RSV/GL2 targeted vascular internal organs: chiefly the spleen and the lung, with the kidney, liver, and muscle also showing some activity. The upper graph demonstrates that the intra-arterial injection of identical amounts of the same luciferase reporter virus resulted in nearly exclusive gene targeting to the heart. The next most active organ was the liver, which had approximately 1/30th the luciferase activity of the heart. Nevertheless, the luciferase activity in the liver was approximately three times higher than that obtained in the spleen or lung following IP injection. The middle graph of Figure 3 shows the effect of employing a muscle-specific promoter to drive the luciferase reporter gene. A 394 bp fragment of the chicken skeletal α -actin promoter is positioned in front of the luciferase reporter gene in this adenoviral vector designated Ad5/sk394/Luc. The results of injecting equivalent titers of this recombinant adenovirus into the carotid artery demonstrate cardiac specificity, but the absolute levels of expression obtained in the heart are 400-500 fold lower than that obtained using the RSV long terminal repeat.

In summary, Figure 3 demonstrates that the direct intra-arterial administration of recombinant gene transfer vectors can target the heart at high efficiency, and intraperitoneal administration can target the spleen and lung with some degree of specificity but at lower efficiency. Previous work from other laboratories demonstrates that the liver can be

-55-

targeted at high efficiency using the same promoter (the RSV LTR), simply by injecting the vector into the portal vein. When combined with the Examples 1-6 demonstrating direct gene transfer into coronary arteries, it is
5 evident that most of the tissues with relevance to the cardiovascular system can specifically be targeted via percutaneous catheterization or by simple intravascular injection. These results demonstrate that somatic gene therapy approaches to cardiovascular diseases can
10 potentially be implemented in a clinical setting using conventional materials and methods.

Example 8

Construction and Evaluation of Synthetic Vectors in the Treatment of Restenosis

15 The idea of using somatic cell gene therapy to inhibit coronary restenosis is appealing because: i) the transferred genes can express the recombinant protein within the restenotic lesion, thus reducing adverse systemic effects, ii) recombinant gene
20 expression may be necessary for only the first few weeks after angioplasty since SMC proliferation is largely confined to this period and iii) a constitutively active gene may be adequate to suppress the restenotic process since precise regulation of gene
25 expression may not be required.

However, the potential of gene therapy to inhibit restenosis has been severely limited by the low efficiency of gene transfer mediated by conventional methods, such as Lipofection. The inefficiency of
30 Lipofection is attributed primarily to the lysosomal destruction of transferred DNA Ho, S. et al., J. Histochem. Cytochem. 31:404-420 (1983).

Coronary smooth muscle cell (SMC) culture was used as an in vitro model of the post-angioplasty vessel wall
35 because few endothelial cells remain in the target

-56-

coronary segment following the angioplasty of atherosclerotic arteries, and because the proliferative response of SMCs to high serum culture conditions in vitro resembles their hyperplastic response to vessel injury in vivo. The species-dependent variability in the susceptibility of cultured SMCs to transfection prompted a comparison of human coronary SMCs with those derived from the coronary arteries of dogs and pigs, the two large animals most commonly used to create models of cardiovascular disease.

Thus, in order to enhance transfection efficiency, cultured human, canine and porcine coronary SMCs were evaluated in two alternative methods of gene transfer. In the first, a synthetic vector was produced in which the hemagglutinin (HA) spike protein of influenza virus was added to the above-described reporter plasmid-Lipofectin™ complexes. The HA spike protein was added to test whether its membrane fusing properties (White, J. et al., Nature 300:658-659 (1982)) would mediate lysosomal escape and improve gene transfer efficiency. In the second method of gene transfer, a recombinant adenoviral vector was employed to directly transfer the reporter gene into SMCs. Although the natural target of the adenovirus is the respiratory epithelium (Rosenfeld, M.A. et al., Cell 68:143-155 (1992)), it has a wide host cell range (Lemarchand, P. et al. Circ. Res. 72:1132-1138 (1993); Salle, G.L.G.L. et al., Science 259:988-990 (1993); Quantin, B. et al., Proc. Natl. Acad. Sci. (U.S.A.) 89:2581-2584 (1992)) and is shown here to efficiently infect cultured SMCs. In addition, the adenovirus has been shown to possess a nuclear transport system that can further enhance gene transfer and expression.

Cell Culture

The human coronary SMCs were obtained from the recipient's explanted heart at the time of cardiac

-57-

transplantation. The porcine coronary SMCs (Campbell, J.H. et al., In: Vascular Smooth Muscle in Culture, Campbell, J.H. et al., (eds.), Vol.1. CRC Press, 1987, pp 15-21)) were harvested from domestic swine and the
5 canine (Feldes, T.F. et al. Amer. J. Physiol. 264:C169-C178 (1993)) from mongrel dogs. The coronary arteries were dissected free of overlying adventitia and minced into small pieces approximately 1 mm square. Primary
10 cultures were established by digesting with an enzyme mixture consisting of 1 mg/ml collagenase (268 U/mg, Worthington Biochemical, Freehold, NJ), 0.25 mg/ml elastase (Grade II, 3 U/mg, BMB, Indianapolis, IN), and 1 mg/ml of soybean trypsin inhibitor (Sigma, St. Louis, MO) in Hanks' balanced salt solution (HBSS). The cells
15 were subsequently grown in Dulbecco's modified Eagle medium (DMEM) with 10% fetal bovine serum (FBS), 0.29 mg/ml L-glutamine, 100 U/ml Penicillin G, 100 µg/ml Streptomycin, and 25 ng/ml Amphotercin (all from GIBCO). The cultures exhibited the characteristic growth pattern
20 (hills and valleys) of SMCs and their identity was confirmed by immunostaining with a monoclonal antibody directed exclusively against smooth muscle α-actin (Sigma). Transfections and infections were performed on SMCs between passages 3 and 5.

25 Transfection

The coronary SMCs were plated 24 hours prior to transfection in 35 mm dishes at a cell density of 5×10^4 cells/cm². At the time of transfection the cells were approximately 80% confluent. Transfection was performed
30 with 1.67 µg of reporter plasmid pRSVL complexed with 5 µg of Lipofectin™ reagent, DOTMA (BRL; Gaithersburg, MD) per dish diluted in 2 ml of OptiMEM I (BRL) for 24 h. The media was subsequently changed to DMEM supplemented with 10% FBS and the cells incubated for an additional
35 48 h. Influenza hemagglutinin (HA) spike protein was obtained from the influenza virus vaccine, trivalent

-58-

types A and B (Wyeth Lab., Marietta, PA). This vaccine of purified subvirion consists of 15 μ g of A/Taiwan, 15 μ g of B/Panama, 15 μ g of A/Beijing in a volume of 0.5 ml. The vaccine stabilizer (thimerosal) was removed and the purified subvirion concentrated by multiple passages through a Centriprep-30 (Amicon Inc., Beverly, MA) to achieve a final concentration of 200 μ g/ml as assessed by the Bio-Rad protein assay (Bio-Rad Laboratories, Richmond, CA). The HA spike protein in amounts of 200, 400, 800 and 1000 ng was added directly to the Lipofectin™/DNA complexes prior to dilution with OptiMEM I and transfection of SMC cultures.

Adenoviral Vector Construction, Propagation, and Infection

The plasmids pFG140 and pJM17 (Frank L. Graham, McMaster University, Ontario, Canada) are derived from the adenovirus type 5 (Ad5) dl 309 genome which has a deletion in the E3 region from map units 83.5 to 85 (Graham, F.L. *et al.*, Meth. Molec. Biol. 7:109-128 (1991)). A 2.2 kb plasmid (pMX2) carrying a plasmid origin of replication and an ampicillin resistance gene has been inserted at the XbaI site in the E1 region of a circularized dl 309 genome to produce the 38.1 kb plasmid, pFG140 (Ghosh-Choudhury, G. *et al.*, Gene 50:161-171 (1986)). The replacement of the 2.2 kb pMX2 plasmid backbone in pFG140 with a larger (4.4 kb) pBR322 derivative results in pJM17, a 40.3 kb modification of the Ad5 dl309 genome which is too large to be packaged into the adenoviral capsid (Ghosh-Choudhury, G. *et al.*, Gene 50:161-171 (1986); McGrory, W.J. *et al.*, Virol. 163:614-617 (1988)). The plasmid pXCJL.1 contains segments from the left-end of the Ad5 genome, from map units 0.0-1.3 and 9.2-15.8 (Frank L. Graham, McMaster University, Ontario, Canada). A 400 base pair (bp) segment of Rous sarcoma virus (RSV) long terminal repeat (LTR), a 1649 bp firefly luciferase cDNA, and 916 bp of

-59-

SV40 sequences including the small-t intron/polyadenylation signal were inserted into pXCJL.1 between Ad5 map units 0.0-1.3 and 9.8-15.8 to generate pXCJL.1/RSV/GL2.

5 The recombinant replication-deficient adenovirus carrying the luciferase cDNA situated behind the RSV LTR (Ad5/RSV/GL2) was obtained through homologous recombination in 293 cells co-transfected with the two plasmids: pJM17 and pXCJL.1/RSV/GL2. The plasmid pJM17
10 alone, after transfection into 293 cells, is incapable of producing a virus because the size constraints of the adenoviral capsid prevent its 40.3 kb genome from being packaged. This promotes homologous recombination
15 between pJM17 and pXCJL.1/RSV/GL2 by selecting for the shorter recombinant which can be rescued into the adenoviral capsid. The resulting virus, Ad5/RSV/GL2 has a substitution of the luciferase expression cassette for the adenoviral E1 genes. The deletion of the E1 genes renders the adenovirus replication deficient in all
20 cells other than the permissive 293 host which provides the E1 functions in trans. Ad5/FG140, on the other hand, was obtained by transfecting 293 cells with plasmid pFG140 alone, since its 38.1 kb can be packaged into the viral capsid. The β -galactosidase reporter
25 virus (Ad5/HCMV/LacZ) was provided by F. L. Graham and A. Bett. This Ad5 derivative contains the human cytomegalovirus (CMV) IE promoter, E. coli lacZ gene, and the SV40 polyadenylation signal inserted in place of the E1 region of the adenoviral genome. The
30 Ad5/RSV/GL2, Ad5/HCMV/LacZ and Ad5/FG140 viruses were plaque-purified and propagated in 293 cells according to published protocols (Graham, F.L. et al., Meth. Molec. Biol. 7:109-128 (1991)) to produce viral stocks with titers of $\geq 2 \times 10^9$ pfu/ml.

35 Infection of porcine, canine and human coronary smooth muscle cells with the replication-deficient adenovirus was performed in 35 mm dishes approaching a

-60-

confluency of 80%. Serial dilutions of the recombinant adenoviruses were prepared in OptiMEM I and cells were exposed to 250 μ l of the virus in OptiMEM I for 30 minutes. The titers of Ad5/RSV/GL2 used for infection were 2.5×10^6 , 1×10^7 , 5×10^7 and 2.5×10^8 pfu/plate. Infection with Ad5/HCMV/LacZ was carried out with viral titers of 5×10^6 , 2.5×10^7 and 1×10^8 pfu/plate. The control dishes were exposed to 250 μ l of Ad5/FG140 in OptiMEM I containing 2.0×10^8 pfu. After 30 minutes of exposure to the adenovirus, 1.75 ml of OptiMEM I was added to the culture plates. The medium was changed at 24 h to DMEM with 10% FBS.

In order to evaluate the effect of exposure time on infection efficiency, porcine SMCs were exposed to Ad5/RSV/GL2 for increasing periods of time. 250 μ l of Ad5/RSV/GL2 (2.5×10^8 pfu) were added to the SMC cultures 24 h after plating at a density of 5×10^4 /cm² in 35 mm culture dishes for exposure times ranging from 30 seconds to 30 minutes. At the end of each interval, the Ad5/RSV/GL2 was aspirated, the plates washed twice with HBSS, and DMEM with 10% FBS was added.

Luciferase Assay

Luciferase activity was assayed seventy-two hours after transfection or infection. Briefly, the SMC cultures were washed with PBS, released with 150 μ l of lysis buffer (25 mM Tris-phosphate, 2 mM dithiothreitol (DTT), 2 mM EDTA, 10% glycerol and 0.1% Triton X-100) and transferred to microfuge tubes. The cell lysate was centrifuged at 12,000 g for 5 minutes, and duplicate enzymatic assays for luciferase activity were performed with 20 μ l samples of supernatant using a Monolight 2010 luminometer (Analytical Luminescence Laboratory [ALL]; San Diego, CA). A 100 μ l aliquot of assay reagent (20 mM tricine, pH 7.8, 1.07 mM (MgCO₃)₄Mg(OH)₂·5H₂O, 2.67 mM MgSO₄, 0.1 mM EDTA, 33.3 mM DTT, 270 μ M coenzyme A, 470 μ M D-luciferin and 530 μ M ATP) was added to each 20

-61-

5 μ l sample, and the microprocessor-controlled photon counter integrated light production over a period of 30 seconds. The luminometer was calibrated using parallel reactions performed with certified firefly luciferase control standard (ALL). Background values obtained from control culture plates (equivalent to background obtained from lysis reagent) were subtracted and values were reported as the total luciferase activity recovered per 35 mm plate.

10 Histochemical Assay for β -galactosidase

The proportion of cells infected with various titers of Ad5/HCMV/LacZ was assessed quantitatively at 72 h by histochemical staining of SMCs for β -galactosidase activity. The SMCs were fixed in 0.5% glutaraldehyde-phosphate buffered saline, pH 7.2, for 5 min at 37°C, then stained with 5 mM $K_4Fe(CN)_6$, 5 mM $K_3Fe(CN)_6$, 2 mM $MgCl_2$, 1 mg/ml 5-bromo-4-chloro-3-indolyl- β -D galactopyranoside (X-gal; BMB) in phosphate buffered saline, pH 7.2, for 16 hr at 37°C. The proportion of cells that expressed β -galactosidase was determined by counting the number of cells stained blue in a representative sample of \approx 200 SMCs per culture plate.

Results of Lipofection With and Without HA

25 The efficiency of gene transfer was measured by quantitating reporter gene protein per 35 mm plate by means of a luminometric assay for luciferase. Using Lipofection alone, the transfection of 1.67 μ g of pRSVL produced 0.016, 2.2 and 5.2 picograms (pg) of reporter protein in human, porcine and canine SMCs, respectively. Thus the efficiency of transfection was quite low, particularly in SMCs of human origin.

In contrast, when increasing amounts of influenza HA spike protein was added to the DNA/Lipofectin™ complexes, a peak stimulation of reporter gene

35

-62-

expression in human and porcine SMCs was obtained with 600 ng of HA per plate. The mean amounts of luciferase protein were 6.9 and 20.3 pg for human and porcine SMCs, respectively, representing approximately a 631- and 9- fold augmentation in gene product over Lipofectin™ alone. In contrast, canine SMCs were far more responsive to Lipofectin™ alone (yielding 5.2 pg/plate) and demonstrated a nearly linear dose-response relationship to HA, yielding 35.7 pg/plate at the highest dose tested (1000 ng).

Species differences in the toxicity of HA on SMCs were also observed. When ≥ 800 ng HA was added per plate, cell death with consequent reduction in gene product was observed in porcine and human SMCs. At 800 ng of HA per plate, toxicity was manifested as cell death in approximately 30 % of these cell populations. Cell death reached 70 % with HA levels of 1000 ng/35 mm dish. The canine SMCs, in contrast, appeared to be unaffected by HA; even when up to 1000 ng of HA was added to a 35 mm culture plate.

Gene Transfer with Recombinant Adenovirus

The proportions of β -galactosidase positive cells in SMC cultures of human, porcine and canine origin were determined by histochemical staining with X-gal chromogen at 72 h after infection with Ad5/HCMV/LacZ. An infection rate of $\geq 95\%$ was evidenced by the presence of blue chromogen in SMCs of all three species at doses of 5×10^6 pfu/35 mm plate. No evidence of β -galactosidase activity was observed in parallel experiments performed with uninfected or Ad5/FG140 (2×10^8 pfu) infected SMC cultures after the 16 h staining period. This demonstrates that the presence of blue chromogen was due to reporter gene expression and not to endogenous β -galactosidase activity, induced either by the cell isolation, culture techniques, or adenoviral infection.

-63-

The Ad5/HCMV/LacZ dose of 1×10^8 pfu/plate was cytotoxic for human and porcine SMCs and resulted in 80% cell death. However, no toxic effects were observed in canine SMCs at similar titers of Ad5/HCMV/LacZ. Cell death was not observed in SMCs of all three species infected with similar doses of Ad5/RSV/GL2.

A quantitative analysis of reporter gene product was performed by a luminometric assay for luciferase in SMCs of all species infected with Ad5/RSV/GL2. The coronary SMCs were exposed to increasing titers of Ad5/RSV/GL2 and assayed for expression of luciferase after 72 h in culture. At low doses of virus (2.5×10^6 pfu/plate), the human and porcine SMCs had similar levels of luciferase; 17 and 12 pg, respectively. However, 7- to 10- fold higher levels of reporter gene product (126 pg) were observed in canine SMCs at the same dose. As the dose was increased to 1×10^7 pfu/plate, species differences in gene expression were less significant. At the highest titer utilized (2.5×10^8 pfu/plate), the quantity of gene product per 35 mm culture plate was 1106, 993 and 1215 pg for human, porcine and canine coronary SMCs, respectively. These values are approximately 113,000-, 450- and 230- fold better than the best results obtained with Lipofectin™ alone in human, porcine and canine coronary SMCs, respectively. Thus the SMCs from all 3 species demonstrated a similar dose-response relation to adenoviral infection, suggesting that the two large animal models may well be appropriate for evaluating gene therapy protocols for inhibiting coronary restenosis.

The duration of exposure to gene transfer reagent is likely to be an important variable in intracoronary gene transfer in vivo. A transfection agent or viral vector that requires short exposure time is desirable because coronary blood flow cannot be interrupted for more than a few minutes during in vivo gene transfer.

-64-

In order to evaluate this parameter, porcine SMCs were exposed for different periods of time to Ad5/RSV/GL2. An exposure duration of only 30 seconds resulted in 18 pg of luciferase protein per 35 mm plate by the
5 luminometric assay. There was no significant difference in reporter gene product resulting from exposure times of 30 sec to 5 minutes. However, when exposure was prolonged to 15 minutes, the level of reporter gene product increased by approximately 3-fold to 43 pg.
10 When compared with Lipofectin™-mediated gene transfer, a 15 minute exposure to 2.5×10^8 pfu/plate of Ad5/RSV/GL2 resulted in gene expression 20-fold higher than that achieved with a 24 h of exposure to Lipofectin™/pRSVL complexes.

15 The data presented herein demonstrate differences in the efficiencies of gene transfer mediated by Lipofectin™ alone, Lipofectin™ plus HA, and recombinant adenovirus in SMCs of human, porcine and canine origin. The low levels of gene transfer and expression observed
20 with Lipofectin™ alone are probably due to lysosomal degradation of the majority of endocytosed DNA. The addition of influenza hemagglutinin in the form of purified subvirion significantly improved the levels of gene transfer and expression in human, porcine and
25 canine SMCs. This improvement is probably due to the escape of the endocytosed complex carrying the plasmid DNA from the perinuclear endosome prior to lysosomal degradation.

Unless specifically engineered to circumvent
30 lysosomal degradation, the fate of most liposomes and their DNA content is to be endocytosed and destroyed. Several strategies have been developed to bypass this problem, including the reconstitution of the membrane proteins of influenza virus and Sendai virus into the
35 liposomal bilayer (Mannino, R.J. et al., Biotechniques 6:682-691 (1988); Kaneda, Y. et al., Exper. Cell Res. 173:56-69 (1987)). Technically complex methods such as

-65-

detergent dialysis and modified EDTA-chelation procedures have been used to prepare these proteoliposomes (Mannino, R.J. *et al.*, Biotechniques 6:682-691 (1988)). For the HA experiments described
5 here, the addition of influenza virus vaccine containing HA spike protein directly to the DNA/Lipofectin™ complexes provides a simple and effective means of increasing gene transfer efficiency.

Although HA augmented gene transfer when compared
10 with Lipofectin™ alone, the best results (6.9 and 20.3 pg/plate) obtained in porcine and human SMCs were still lower than recombinant adenovirus-mediated gene transfer. Recombinant adenovirus-mediated gene transfer resulted in the synthesis of approximately 1 ng/plate of
15 luciferase in SMC cultures of all three species. Similar levels of expression from a gene encoding an antiproliferative or anticoagulant protein would be expected to have a therapeutic effect on restenotic lesions in vivo. In addition, the ability of adenoviral
20 vectors to mediate gene transfer during relatively short periods of exposure to SMCs is a very desirable feature considering the time constraints on the delivery of in vivo intracoronary gene therapy.

Species differences in the susceptibility of SMCs
25 to transfection and toxicity from Lipofectin™ plus HA and recombinant adenovirus were evident. The canine SMCs were unique among the three species tested because they were readily transfected by Lipofectin™ alone and were more resistant to the cytotoxic effects of HA or
30 recombinant adenovirus. The similarity between human and porcine SMCs in their response to these two gene transfer methods suggests that swine may be a more suitable model in which to evaluate potential transfection agents or viral vectors for the in vivo
35 delivery of gene therapy.

In summary, in vitro studies with coronary SMCs

-66-

from three species suggest that adenoviral-mediated gene transfer is highly efficient. By comparison, the technique of Lipofection (even when enhanced with HA spike protein) results in relatively low levels of gene transfer and expression. Although HA significantly enhanced the efficiency of Lipofection in coronary SMCs from all three species, the maximal enhancement was often dose-dependent. Canine cells were unique in their resistance to HA toxicity, while human and porcine cells exhibited a sensitivity to HA which resulted in cell death and decreased reporter gene expression at higher doses. Similarly, canine SMCs were resistant while human and porcine cells were sensitive to high doses of the recombinant adenovirus expressing β -galactosidase. In addition to being far more efficient than Lipofectin™-based methods, the adenovirus expressing luciferase produced similar amounts of recombinant protein in SMCs from all three species. Thus, the results of adenovirus-mediated gene therapy strategies to prevent restenosis in canine and particularly porcine models appear fully applicable to humans. The present invention demonstrates that adenovirus is an efficient vector for gene transfer into cultured SMCs and a particularly attractive candidate for the in vivo delivery of therapeutic genes into coronary arteries following angioplasty.

While the invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications and this application is intended to cover any variations, uses, or adaptations of the invention following, in general, the principles of the invention and including such departures from the present disclosure as come within known or customary practice within the art to which the invention pertains and as may be applied to the essential features hereinbefore

-67-

set forth and as follows in the scope of the appended claims.

-68-

WHAT IS CLAIMED IS:

1. A method of genetic therapy for the treatment of cardiovascular disease, wherein said method comprises providing an effective amount of a gene transfer vector to a somatic cell of an animal, said vector comprising a therapeutic gene sequence and being sufficient to mediate efficient gene transfer.
2. The method of claim 1, wherein said gene transfer vector is derived from adenovirus subgenus C.
3. The method of claim 1, wherein said gene transfer vector is a complex containing two or more of the following components or their synthetic or carbohydrate derivatives: protein, cationic polymer, or cationic lipid.
4. The method of claim 3, wherein a component of said vector provides at least one function selected from the group consisting of: i) plasma membrane penetration; ii) escape from lysosomal degradation; iii) nuclear transport; and iv) persistent nuclear retention.
5. The method of claim 4, wherein said plasma membrane penetration is mediated by membrane fusion or endocytosis.
6. The method of claim 4, wherein said escape from lysosomal degradation is mediated by endosomal avoidance or disruption.
7. The method of claim 4, wherein said nuclear transport is mediated by a nuclear localization sequence.

-69-

8. The method of claim 4, wherein said persistent nuclear retention is by mediated by resistance to nuclease activity or by providing for replication.

5 9. The method of claim 3, wherein said complex contains cationic polymers or cationic lipids, and wherein the structure of said complex is maintained by ionic interactions between said nucleic acid of said vector and said cationic polymers or cationic lipids.

10 10. The method of claim 3, wherein said complex contains protein and a cationic polymer, and wherein said protein is conjugated to said cationic polymer, and wherein the structure of said complex is maintained by ionic interactions between said nucleic acid of said vector and said conjugated protein and cationic polymer.

15 11. The method of claim 3, wherein said complex contains protein and wherein said protein is incorporated into a liposomal complex of cationic lipids, said complex being in association with said nucleic acid of said vector.

20 12. The method of claim 1, wherein said animal is a human.

13. The method of claim 1, wherein said somatic cell is a smooth muscle cell.

25 14. The method of claim 1, wherein said somatic cell is an endothelial cell.

15. The method of claim 1, wherein said somatic cell is a cardiomyocyte.

16. The method of claim 1, wherein said somatic cell is a cardiac fibroblast.

-70-

17. The method of claim 1, wherein said somatic cell is a cell of the immune system.

18. The method of claim 1, wherein said somatic cell is a hepatocyte.

5 19. The method of claim 1, wherein said cardiovascular disease is restenosis.

20. The method of claim 1, wherein said cardiovascular disease is atherosclerosis.

10 21. The method of claim 1, wherein said cardiovascular disease is dyslipidemia.

22. The method of claim 1, wherein said cardiovascular disease involves inflammation.

23. The method of claim 1, wherein said cardiovascular disease involves thrombosis.

15 24. The method of claim 1, wherein said cardiovascular disease involves ischemia.

25. The method of claim 1, wherein said cardiovascular disease involves myocardial infarction.

20 26. The method of claim 1, wherein said cardiovascular disease involves reperfusion injury.

27. The method of claim 1, wherein said cardiovascular disease involves immunological rejection of a transplanted organ or tissue.

25 28. The method of claim 1, wherein said therapeutic gene sequence is expressed by said somatic cell.

-71-

29. The method of claim 28, wherein said therapeutic gene sequence encodes a protein selected from the group consisting of thrombin inhibitors, platelet inhibitors, GF inhibitors, antioxidants, antiproliferative proteins, chimeric toxins, anti-inflammatory agents, antilipemic agents, anti-thrombotics, transplant rejection inhibitors and a soluble receptor.

30. The method of claim 28, wherein said protein is arrayed on a cell surface of said somatic cell.

31. The method of claim 30, wherein said arrayed protein is selected from the group consisting of a receptor, a receptor antagonist, and a dominant-negative mutant receptor protein.

32. The method of claim 1, wherein said therapeutic gene sequence encodes an antisense or ribozyme RNA.

33. The method of claim 1, wherein said method is conducted in conjunction with an interventional or surgical procedure.

34. The method of claim 1, wherein said effective amount of said vector is provided in vivo by a percutaneous, catheter-based approach.

35. The method of claim 1, wherein said effective amount of said vector is provided in vivo by direct intra-arterial injection or infusion, with or without surgical isolation of said vessel.

36. The method of claim 1, wherein said effective amount of said vector is provided in vivo by direct intravenous injection or infusion.

37. The method of claim 1, wherein said effective amount of vector is provided ex vivo to a graft, tissue, organ or cell population, and wherein said graft, tissue, organ or cell population is then introduced into said animal.

38. The method of claim 1, wherein the expression of therapeutic gene sequences is controlled by a tissue-specific promoter, a developmentally regulated promoter or an enhancer element.

39. The method of claim 1, wherein the expression of said therapeutic gene sequence is controlled by a promoter, or an enhancer element, and wherein said control is responsive to the systemic administration of an exogenous substance.

40. The method of claim 39, wherein a gene sequence encoding a recombinant receptor is also administered, and wherein said recombinant receptor provides a signal transduction for said exogenous substance.

41. The method of claim 1, wherein said method comprises providing more than one therapeutic gene sequence to said somatic cell of said animal.

42. The method of claim 41, wherein said vector comprises said more than one therapeutic gene sequence.

43. The method of claim 41, wherein said therapeutic gene sequences are provided to said somatic cell of said animal by a plurality of vectors.

44. The method of claim 41, wherein said therapeutic gene sequences are provided to said somatic cell in conjunction with pharmacological therapies.




FIG. 1

SUBSTITUTE SHEET (RULE 26)

2/3




FIG. 2

SUBSTITUTE SHEET (RULE 26)




FIG. 3A




FIG. 3B




FIG. 3C

SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/05548

A CLASSIFICATION OF SUBJECT MATTER

CLASS. A61K 31/675; C12N 5/00, 15/00

US CL. 514/44; 435/69.1, 172.1, 172.3, 320.1

According to International Patent Classification (IPC) or to both national classification and IPC

B FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

US 514/44; 435/69.1, 172.1, 172.3, 320.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	THE JOURNAL OF REPRODUCTIVE MEDICINE, Volume 37, Number 6, issued June 1992, E.M. Karson et al., "Prospects for Human Gene Therapy", pages 508-514, see entire document.	1-44
Y	HUMAN GENE THERAPY, Volume 2, issued 1991, F.D. Ledley, "Clinical considerations in the design of protocols for somatic gene therapy", pages 77-83, see entire document.	1-44
Y	CHEMICAL REVIEWS, Volume 90, Number 4, issued June 1990, E. Uhlmann et al., "Antisense oligonucleotides: A new therapeutic principle", pages 543-584, see entire document.	1-44

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

A	document defining the general state of the art which is not considered to be of particular relevance	*T*	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
E	earlier document published on or after the international filing date	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
L	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
O	document referring to an oral disclosure, use, exhibition or other means	*Z*	document member of the same patent family
P	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

21 JULY 1994

Date of mailing of the international search report

09 AUG 1994

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

BRIAN R. STANTON

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/05548

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	CELL, Volume 68, issued 10 January 1992, M.A. Rosenfeld et al., "In vivo transfer of the human cystic fibrosis transmembrane conductance regulator gene to the airway epithelium", pages 143-155, see entire document.	1-44
Y	SCIENCE, Volume 249, issued 14 September 1990, E.G. Nabel et al., "Site-specific gene expression in vivo by direct gene transfer into the arterial wall", pages 1285-1288, see entire document.	1-44
Y	CIRCULATION, Volume 85, Number 3, issued March 1992, M.Y. Flugelman et al., "Low level in vivo gene transfer into the arterial wall through a perforated balloon catheter", pages 1110-1117, see entire document.	1-44

Form PCT/ISA/210 (continuation of second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US94/05548

B FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

APS, MEDLINE, CA

search terms: NABAL?/AU; FRENCH?/AU; ROBERTS?/AU; GENE?; THERAPY; CARDIOVASCUL?; RESTENO?;
ATHEROSCLERO?; INFLAMM?; THROMBOSIS; ISCHEMI?; REPERFUSION; INJUR?; ADENOVIR?

Form PCT/ISA/210 (extra sheet)(July 1992)*